

doi.org/10.3114/fuse.2021.07.09

Taxonomy and phylogeny of the basidiomycetous hyphomycete genus *Hormomyces*

J. Mack*, R.A. Assabgui, K.A. Seifert#

Biodiversity (Mycology and Microbiology), Agriculture and Agri-Food Canada, 960 Carling Avenue, Ottawa, Ontario K1A 0C6, Canada. #Current address: Department of Biology, Carleton University, 1125 Colonel By Drive, Ottawa, Ontario K1S 5B6, Canada.

*Corresponding author: jonathan.mack2@canada.ca

Key words:

Dacrymyces
Oosporidium
Tremella
Tulasnella
1 new taxon

Abstract: The taxonomy of the genus *Hormomyces*, typified by *Hormomyces aurantiacus* which based on circumstantial evidence was long assumed to be the hyphomycetous asexual morph of *Tremella mesenterica* (*Tremellales*, *Tremellomycetes*) or occasionally *Dacrymyces* (*Dacrymycetales*, *Dacrymycetes*), is revised. Phylogenies based on the three nuclear DNA markers [internal transcribed spacers (ITS), 28S large ribosomal subunit rDNA (28S) and 18S small ribosomal subunit rDNA (18S)], based on cultures from Canada and the United States, suggest that the genus is synonymous with *Tulasnella* (*Cantharellales*, *Agaricomycetes*) rather than *Tremella* or *Dacrymyces*. Morphological studies of 38 fungarium specimens of *Hormomyces*, including the type specimens of *H. callorioides*, *H. fragiformis*, *H. paridiphilus* and *H. peniophorae* and examination of the protologues of *H. abieticola*, *H. aurantiacus* and *H. pezizoideus* suggest that *H. callorioides* and *H. fragiformis* are conspecific with *H. aurantiacus* while the remaining species are unlikely to be related to *Tulasnella*. The conidial chains produced by *H. aurantiacus* are similar to monilioid cells of asexual morphs of *Tulasnella* species formerly referred to the genus *Epulorhiza*. The new combination *Tulasnella aurantiaca* is proposed and the species is redescribed, illustrated and compared with similar fungi. The ecological niche of *T. aurantiaca* and its possible relationship to orchid root endophytes is discussed. A key to asexual genera with similar conidium ontogeny to *T. aurantiaca* is provided.

Citation: Mack J, Assabgui RA, Seifert KA (2021). Taxonomy and phylogeny of the basidiomycetous hyphomycete genus *Hormomyces*. *Fungal Systematics and Evolution* 7: 177–196. doi: 10.3114/fuse.2021.07.09

Received: 20 November 2020; Accepted: 21 January 2021; Effectively published online: 12 February 2021

Corresponding editor: P.W. Crous

INTRODUCTION

The genus *Hormomyces* was introduced by Bonorden (1851) for a single hyphomycetous species, *H. aurantiacus* collected on old oak (*Quercus* sp.) wood in Germany. The protologue described and illustrated orange, gelatinous sporodochia with branched chains of hyaline, globose conidia; no measurements for microscopic structures were reported (Fig. 1). The location of Bonorden's herbarium is uncertain (Stafeu & Cowan 1976) and no type specimen is known to exist for this species. *Hormomyces*




Fig. 1. A copy of the lectotype figure for *Hormomyces aurantiacus*, a reproduction of fig. 234, taf. XI from Bonorden (1851).

is rarely mentioned in modern taxonomic literature (Seifert *et al.* 2011). The only modern study is by Tubaki (1976), who reported on its blastoacropetal conidium ontogeny and the characters of *H. aurantiacus* in axenic culture based on a strain isolated from a fallen twig of *Tsuga canadensis* collected in New York State, United States. His concept was generally accepted. The colonies of *H. aurantiacus* are conspicuous and distinctive but they may be disregarded as uninteresting “jelly fungi” by those focusing on microfungi, or discarded as a “trivial asexual morph or anamorph” by those interested in macrobasidiomycetes. Despite this lack of academic attention, there are frequent reports by field mycologists, mostly accurately reflecting the Tubaki (1976) concept, e.g. Mushroom Observer includes 42 (Wilson *et al.* 2020) and MyCoPortal 180 records (MyCoPortal 2019).

Seven species of *Hormomyces* were described: *H. abieticola*, *H. aurantiacus*, *H. callorioides*, *H. fragiformis*, *H. paridiphilus*, *H. pezizoideus* and *H. peniophorae* (supplementary table S1). Two were first described in *Hypsilophora* prior to being transferred to *Hormomyces* by Saccardo (1888), namely *Hy. fragiformis* and *Hy. callorioides*. Saccardo (1888), who mostly compiled descriptions from other mycologists and often did not examine material himself, considered the main distinction among these species to be sporodochial colour: orange for *H. aurantiacus*, purple for *H. fragiformis* and pink for *H. callorioides*. Lloyd (1916) questioned the value of colour to distinguish these species, suggesting that they might be conspecific without formally synonymizing them.

McNabb (1969) synonymized *H. callorioides* under *H. fragiformis*, re-emphasizing their exclusion from *Hypsiloophora*, which he considered the correct generic name for the asexual morph of the fungus now known as *Erythricium salmonicolor* (*Corticiales*). McNabb did not propose a synonymy of *H. fragiformis* with *H. aurantacus*, and the former name has often been used for specimens collected in North America (MyCoPortal 2019), irrespective of sporodochial colour. Tubaki (1976) accepted only *H. aurantacus* in the genus, although he did not critically revise the other named species.

In their compilation of hyphomycete genera, Seifert *et al.* (2011) listed two other genera as synonyms of *Hormomyces*: *Sphaerocolla* (following the opinions of von Höhnelt 1917; Donk 1962) and *Hormisciopsis*. *Sphaerocolla* Karsten (1892) comprises *S. aurantaca* from Finland from living *Betula* bark, reported as producing large, effuse, orange sporodochia up to 10 cm long, with branched chains of globose conidia 3–9 µm diam, reminiscent of *H. fragiformis* or *H. aurantacus*. *Hormisciopsis* Sumstine (1914) was proposed for *Ho. gelatinosa*, producing red, gelatinous sporodochia on wood collected in Pennsylvania, with branching chains of globose to ellipsoidal conidia 6–10 × 5–6 µm. The description corresponds well with *H. fragiformis* and *H. aurantacus*, and the illustration is very similar to those of *H. aurantacus* provided by Lloyd (1916) and *H. fragiformis* by Patouillard (1900). Although Sumstine (1914) compared *Hormisciopsis gelatinosa* to *Hormiscium*, he was apparently unaware of *Hormomyces*.

The taxonomic relationships of *Hormomyces* were a matter of debate, although it has usually been regarded as a basidiomycete. Patouillard (1900) suggested that *Hormomyces* is the budding state of a *Dacrymyces*. Lloyd (1916) disagreed but did not provide an alternative classification. That same year, Saccardo suggested that *H. aurantacus* might be the asexual morph of *Tremella mesenterica* (Saccardo 1916). Bresadola (1932) reiterated this putative connection in a short note, reporting that the fungus he examined had conidia up to 3 µm long but he did not illustrate it; unfortunately, no specimens were mentioned. Donk (1962) also suggested a relationship between *Tr. mesenterica* and *H. aurantacus*, noting that immature *Tr. mesenterica* also produced conidia. McNabb (1969) suggested that this connection was “currently accepted in Europe.” Later, however, Pippola & Kotiranta (2008) described the asexual morph of *Tr. mesenterica* as producing chains of ellipsoidal conidia (1.8–)2.2–4.5(–5.9) × (1.6–)1.8–3.8(–4.2) µm, originating from clamped conidiogenous cells, which matches the dimensions given by Bresadola (1932). The characters diverge significantly from *H. aurantacus* as described by Bonorden (1851), *i.e.* with globose conidia and no clamp connections. Tubaki (1976) also disagreed with this supposed connection, remarking that many asexual morphs of *Tremella* species are yeasts, while his culture of *H. aurantacus* was filamentous. Thus, the purported asexual–sexual morph connection was a speculation that was never established experimentally and was contradicted by subsequent observations. Given their different ecological niches, *H. aurantacus* on rotten wood and *Tr. mesenterica* as a mycoparasite usually on rather solid wood, it is unlikely that the two were seen in close proximity.

In 2015, we isolated a culture from a freshly collected specimen of *H. aurantacus* and our initial 18S rDNA sequences obtained from that culture suggested an affinity of *Hormomyces* with *Tulasnella* rather than *Tremella* or *Dacrymyces*. The genus *Tulasnella* contains

about one hundred sexual species, most of which occur on wood, and most known asexual species are usually isolated from orchid roots or liverworts. The basidiomes are usually thin and waxy and the basidia have four basally swollen sterigmata delimited from the probasidium by a septum. So far, no asexual species of *Tulasnella* have been recorded from wood. The morphology and ontogeny of the conidial chains of *H. aurantacus* are similar to the moniloid hyphae of the orchid mycorrhizal *Tulasnella* asexual morphs formerly attributed to *Epulorhiza* (Roberts 1994), and together with its dikaryotic hyphae, support a relationship with *Tulasnellaceae*. In this paper, our goal is to clarify the family and genus level classifications of this fungus using Bayesian Inference (BI) and Maximum Likelihood (ML) analyses of nuclear DNA regions and genes, and to evaluate the morphological characters of *H. aurantacus* and the other described species of the genus based on studies of types and supplementary specimens and cultures. Should *Hormomyces* be considered distinct from or a synonym of *Tulasnella*? And if *H. aurantacus* is classified in *Tulasnella*, might it be the asexual morph of a known sexual species, or identical with one of the other asexual species previously attributed to *Epulorhiza*? Because *H. aurantacus* is apparently lignicolous, understanding its relationship with other *Tulasnella* species and mycorrhizal species formerly classified in *Epulorhiza* may provide insight into ecological patterns in this family.

MATERIALS AND METHODS

Cultures, specimens and morphological examination

Fresh specimens of *H. aurantacus* were collected in 2014 and 2015 near Ottawa, Ontario, Canada, with additional specimens and one culture provided by colleagues; all material studied is listed in the Specimens Examined paragraphs of the Taxonomy section. Cultures were obtained by squashing a small fragment of a sporodochium and making a slurry that was further diluted with sterile dH₂O. Ten 20 µL drops of the dilution were pipetted onto malt extract agar (MEA, recipe from Raper and Thom 1949), using BD Bacto Malt Extract, and 15 g/L BD Bacto Agar, with the addition of 0.1 g/L ZnSO₄·7H₂O and 0.005 g/L CuSO₄·5H₂O, and examined 24 h later for germinating conidia, which were then transferred individually to new MEA plates to reduce the risk of contamination. Thereafter, we did not attempt to make single spore isolations because the conidia produced *in vivo* remain in strongly coherent chains and can be separated only with difficulty. No evidence of contamination of the cultures by other fungi or bacteria was evident under the dissecting microscope or by light microscopy of mounts from the original isolates.

Culture characters were observed by growing all strains on MEA and oatmeal agar (OA; Gams *et al.* 1987).

For each strain and medium, one plate was incubated under near-UV light for 16 h and in darkness for 8 h per day at ambient RT (varying between 20–25 °C), and another in darkness in a 25 °C incubator. Specimens were inoculated at a single, central point in a 90 mm polystyrene Petri dish. All cultures were then examined and measured every 7 d for 4 wk, with coloration noted on days 14 and 28. To determine cardinal temperatures, two strains were grown in darkness at temperatures ranging from 5–40 °C, at increments of 5 °C (except 37 °C was used instead of 35 °C), and measured every 7 d for 4 wk. Culture photographs were taken with an Olympus Tough TG-5 camera (Olympus, Tokyo) with black velvet as a background.

Herbarium specimens, including available types, were borrowed from BPI (26 specimens), DAOM (7), K(M) (5) and HKAS (1) (herbarium acronyms follow Thiers 2019).

Microscopic observations of living specimens were made using tissue mounted in water. Herbarium specimens were rehydrated with a small drop of dH₂O for 10 min, then small fragments of sporodochia were removed using jewellers' forceps and heated in 85 % lactic acid for five minutes on an electric hot plate. Herbarium specimens and cultures were examined using an Olympus SZX12 dissecting microscope and BX50 compound microscope (Olympus, Tokyo), and photographed using Infinity 2 or Infinity X USB microscope cameras, using Infinity Capture software (Lumenera, Ottawa). Specimen and colony colours were described using the alphanumeric codes and names (with initial capitals) used by Kornerup & Wanscher (1976). Photographic plates were assembled using Photoshop v. 5.5 (Adobe Systems, San Jose, CA).

To demonstrate dikaryotic nuclei, hyphae from a 9-d-old culture of DAOMC 251988 grown on MEA were mounted in 10 µL of SYTO 13 green fluorescent nucleic acid stain (Invitrogen, Carlsbad, California) on a microscope slide. After the placement of the cover glass, the sample was left in darkness for 30 min at 37 °C, then examined with a Nikon ECLIPSE E800 fluorescence microscope (Nikon, Tokyo) using the UV2 setting. Pictures were taken with a Nikon DS-Ri2 camera (Nikon, Tokyo).

For each specimen, 50 conidia were measured, and mean and standard errors are provided. Q values for conidia were calculated as length divided by width. To assess the possible significance of conidial dimensions as a diagnostic character for species identification, means (with upper and lower limits represented by standard error, and outliers in brackets) were compared for all specimens. All calculations were made with Excel 2016 (Microsoft, Redmond).

DNA extraction, sequencing and phylogenetic analysis

Genomic DNA was extracted from pure cultures of *H. aurantacus* using the DNeasy UltraClean Microbial Kit (Qiagen, Hilden), following the manufacturer's protocol. Permission was not obtained to attempt extraction of genomic DNA from specimens loaned by fungaria. For preliminary phylogenetic placement of an early isolate of *H. aurantacus* (DAOMC 252084), a partial sequence of the 18S nrRNA gene was generated using primers NS1 and NS4, and an amplification profile of 95 °C for 10 min for the initial denaturing of the DNA template, then 40 cycles with denaturation at 95 °C for 60 s, annealing at 56 °C for 45 s and extension at 72 °C for 90 s, with a final extension at 72 °C for 10 min. Because of the relationship with *Tulasnellaceae* (*Cantharellales*) suggested by these results, we tried a *Tulasnella*-specific ITS4 primer designed by Taylor & McCormick (2008) in combination with ITS5; this resulted in sporadic and low yield amplification for our strains of *H. aurantacus*. Amplification of the 28S nrRNA gene with LROR and either LR8 or LR5 invariably failed or yielded multiple PCR bands. Amplification was eventually achieved using V9G as the forward primer and LR3 as the reverse primer. See Results for primer references. The PCR profile had an initial denaturation at 94 °C for 90 s, five cycles of denaturation at 94 °C for 45 s, annealing at 56 °C for 45 s and extension at 72 °C for 60 s. Annealing temperatures were then decreased by 1 °C every five cycles until reaching 51 °C, which was used for another 30 cycles, with a final extension of 10 min, for a total of 55 cycles.

Our sequences of *H. aurantacus* were edited and trimmed using Geneious v. 11.1.5 (Biomatters, Auckland). BLAST analyses of all three nrDNA markers indicated a relationship to reference sequences in *Tulasnellaceae*, which guided our sampling for subsequent phylogenetic analyses. Our sequences were aligned using MUSCLE v. 3.8.425 (Edgar 2004) in datasets containing sequences of *Cantharellales* downloaded from GenBank (Table 1 for ITS, Table 2 for 28S), sampled to include all available taxa of *Tulasnellaceae* and a few similar, mostly ITS, environmental sequences and unidentified species uncovered during BLAST searches, along with selected representatives of other genera in the order. We also used an ITS and partial 28S sequence of *H. aurantacus* (NBRC 30400, the culture referred to by Tubaki 1976) published in the NBRC culture collection database (NITE Biological Resource Center 2019), which were not deposited in GenBank. Reference sequences of *Craterellus tubaeformis* were used as the outgroup for the ITS analysis, which examined relationships among species of *Tulasnella*, *Epulorhiza* and *H. aurantacus*. A reference sequence of *Ustilago maydis* was used as the outgroup for the LSU analysis, which focused on the relationship between *Hormomyces* and *Tulasnella* in the *Cantharellales*, and tested prior hypotheses of relationships between *Hormomyces* and *Tremella* or *Dacrymyces*.

Independent phylogenetic analyses were conducted using the Maximum Likelihood (ML) and the Bayesian Inference (BI) algorithms for both markers. The ML analysis was done with PHYML v. 3.0 using the GTR + G + I model as the most suitable model for both 28S and ITS (Guindon *et al.* 2010). For BI, JModelTest v. 0.1.1 (Darriba *et al.* 2012) was used to determine the most suitable model, GTR + G for both ITS and 28S. The BI analyses were run using MrBayes v. 3.2 (Ronquist *et al.* 2012), with four simultaneous Markov chains run until the average standard deviation of split frequency reached < 0.01. Convergence was assessed when the standard deviation of split frequency reached < 0.01. Sampling frequency was 1 in 500 generations with the first 25 % of the trees discarded as burnin. Trees were visualized using FigTree v. 1.4.3 (Rambaut 2016) and modified using Adobe Illustrator 10 (Adobe, San Jose) and PowerPoint 2016 (Microsoft, Redmond). The alignments and phylogenetic trees were deposited in TreeBASE (Treebase.org Study ID: 25624 and 27256). Proposed new names and typifications were deposited in MycoBank (MB) and the MycoBank typification (MBT) database (Westerdijk Fungal Biodiversity Institute, Utrecht).

RESULTS

Phylogenetic analysis

An initial 1045 bp sequence of the 5' end of the 18S nrRNA gene of DAOMC 252084 (deposited as GenBank MN719097) indicated a relationship of *H. aurantacus* with *Tulasnella*, with a 99.2 % similarity to an AFTOL generated sequence of *Tulasnella violea* (AY707097), based on a query coverage of 99 %, with sequence similarities of about 80–99 % with other *Tulasnellaceae*, mostly with query coverages of about 55 %. The distance tree accompanying the BLAST search was consistent with the close placement of our sequence in this family. Because the taxon sampling for 18S sequences of *Tulasnellaceae* is so sparse in GenBank, with only three named species among the 32 reference sequences available, we did not pursue further 18S analyses.

Table 1. ITS sequences retrieved from GenBank or newly generated during this study.

| Species | Strain no. | Origin | Host | GenBank accession no. | Reference |
|------------------------------------|---------------------------|-----------------------------------|---|----------------------------|--|
| <i>Ceratobasidium albasitensis</i> | EaB-T2 | – | – | AJ427398 | Gonzalez, unpublished |
| <i>C. angust sporum</i> | CBS 568.83 | Eyre Peninsula, SA, Australia | <i>Pterostylis mut ca</i> endophyte | AJ427403 | Gonzalez, unpublished |
| <i>C. cereale</i> | C13 | – | <i>Trit cum aest vum</i> | AJ302009 | Gonzalez <i>et al.</i> (2002) |
| <i>C. cornigerum</i> | C6 | – | <i>Erigeron canadensis</i> | AJ301902 | Gonzalez <i>et al.</i> (2002) |
| <i>C. ramicola</i> | CBS 758.79 ^f | Florida, USA | <i>Pit osporum</i> , leaf | AJ427404 | Gonzalez, unpublished |
| <i>C. stevensii</i> | CBS 477.82 | Kentucky, USA | <i>Malus domest ca</i> , twig | AJ427405 | Gonzalez, unpublished |
| <i>Craterellus tubaeformis</i> | S9 | – | – | MH394713 | Jensen-Vargas & Marizzi (2018) |
| | 1D3 | Kunigami, Okinawa, Japan | – | AB973729 | Matsuoka, unpublished |
| <i>Tulasnella albida</i> | KC 110 | – | – | AY373294 | McCormick <i>et al.</i> (2004) |
| <i>T. amonilioides</i> | – | – | – | JF907600 | Almeida <i>et al.</i> (2014) |
| <i>T. anat cula</i> | 13o004 | Mt. Hambeak, South Korea | <i>Platanthera chlorantha</i> | KT164598 | Direct submission |
| <i>T. asymmetrica</i> | MA FF305808 Clone C002 | Australia | <i>Thelymitra epipactoides</i> | KC152347 | Cruz <i>et al.</i> (2014) |
| | MAFF 305808 Clone C005 | Australia | <i>Thelymitra epipactoides</i> | KC152348 | Cruz <i>et al.</i> (2014) |
| | MAFF305808 Clone C009 | Australia | <i>Thelymitra epipactoides</i> | KC152349 | Cruz <i>et al.</i> (2014) |
| <i>T. aurant aca</i> | DAOMC 251988 | Pennsylvania, USA | Rot en wood | MK626686 | This study |
| | DAOM 970795 | | | | |
| | DAOMC 251989 | Tennessee, USA | Rot en wood and <i>Crepidotus</i> spp. | MK626533 | This study |
| | PBM4158 | | | | |
| | DAOMC 252083 | Victoriaville, Quebec, Canada | <i>Fomitopsis betulina</i> | MK626687 | This study |
| | DAOM 970821 | | | | |
| | DAOMC 252084 | Ot awa, Ontario, Canada | Rot en <i>Populus</i> wood | | This study |
| | DAOM 970822 | | | MK626567 | |
| | DAOMC 252086 | Montreal, Quebec, Canada | Rot en wood | MK593626 | This study |
| | DAOM 970819 | | | | |
| | DAOMC 252085 | Ot awa, Ontario, Canada | Rot en wood | MK626568 | This study |
| | DAOM 970820 | | | | |
| | NBRC 30400 | New York, USA | Twig of <i>Tsuga canadensis</i> | Sequence not in Genbank | NITE Biological Resource Center, 2019 |
| <i>T. australiensis</i> | CLM 031 | New York, USA | <i>Arthrochilus oreophilus</i> | KF476602 | Arif n <i>et al.</i> (2020) |
| | CLM 1945 ^f | Nowra, NSW, Australia | <i>Cryptostylis erecta</i> root | MT003730 | Arif n <i>et al.</i> (2020) |
| | CLM 2004 | Northern Sydney NSW, Australia | <i>Cryptostylis erecta</i> root | MT003715 | Arif n <i>et al.</i> (2020) |
| <i>T. calospora</i> | CBS 326.47 | – | – | AY373298 | McCormick <i>et al.</i> (2004) |
| | Ch5-3 | – | – | HM450045 | Idris (2010) |
| <i>T. concentrica</i> | CLM 2071 | Morton NP, NSW, Australia | <i>Cryptostylis leptochila</i> root | MT036547 | Arif n <i>et al.</i> (2020) |
| | CLM 2098 ^f | Nowra, NSW, Australia | <i>Cryptostylis erecta</i> root | MT003744 | Arif n <i>et al.</i> (2020) |
| | CLM2198 | Bunyip SP, Vic, Australia | <i>Cryptostylis leptochila</i> root | MT036533 | Arif n <i>et al.</i> (2020) |
| <i>T. cumulopunt oides</i> | MAFF 245682 | Tsukuba City, Ibaraki, Japan | <i>Spiranthes sinensis</i> | LC175323 | Fujimori <i>et al.</i> (2019) |

Table 1. (Continued).

| Species | Strain no. | Origin | Host | GenBank accession no. | Reference |
|-----------------------------|--------------------------|---|-----------------------------------|-----------------------|--------------------------------|
| <i>T. danica</i> | KC 388 | – | – | AY373297 | McCormick <i>et al.</i> (2004) |
| <i>T. deliquescens</i> | MAFF 244717 | – | <i>Spiranthes sinensis</i> | LC175329 | Fujimori <i>et al.</i> (2019) |
| <i>T. densa</i> | CLM 2110 | Bulahdelah, NSW, Australia | <i>Cryptostylis hunteriana</i> | MT036526 | Arif <i>n et al.</i> (2020) |
| | CLM 2111 | Bulahdelah, NSW, Australia | <i>Cryptostylis hunteriana</i> | MT036525 | Arif <i>n et al.</i> (2020) |
| | CLM 2117 ^f | Bulahdelah, NSW, Australia | <i>Cryptostylis hunteriana</i> | MT036520 | Arif <i>n et al.</i> (2020) |
| <i>T. dentrit ca</i> | MAFF 244709 ^f | – | <i>Spiranthes sinensis</i> | LC175308 | Fujimori <i>et al.</i> (2019) |
| <i>T. eichleriana</i> | KC 852 | – | – | AY373292 | McCormick <i>et al.</i> (2004) |
| | K(M) 143600 | England, UK | Wood of decort cated sapling | KC152381 | Cruz <i>et al.</i> (2014) |
| <i>T. ellipsoidea</i> | MAFF 245686 | Tsukuba City, Ibaraki, Japan | <i>Spiranthes sinensis</i> | LC175315 | Fujimori <i>et al.</i> (2019) |
| <i>T. epiphyt ca</i> | AERO_3.2 | – | – | JF907598 | Almeida <i>et al.</i> (2014) |
| <i>T. irregularis</i> | CBS 574.83 ^f | NT, Australia | <i>Dendrobium dicupum</i> root | MH861654 | Vu <i>et al.</i> (2019) |
| <i>T. occidentalis</i> | CLM 1938 ^f | Boyanup, WA, Australia | <i>Cryptostylis ovata</i> root | MT008096 | Arif <i>n et al.</i> (2020) |
| | CLM 1942 | Boyanup, WA, Australia | <i>Cryptostylis ovata</i> root | MT008092 | Arif <i>n et al.</i> (2020) |
| | CLM 1943 | Boyanup, WA, Australia | <i>Cryptostylis ovata</i> root | MT008091 | Arif <i>n et al.</i> (2020) |
| <i>T. prima</i> | CLM 159 ^f | Blue Mountains, NSW, Australia | <i>Chiloglot s trilabra</i> | KF476556 | Linde <i>et al.</i> (2013) |
| | CLM 377 | Kanangra Boyd NP, NSW, Australia | <i>Chiloglot saf. jeanesii</i> | KF476544 | Linde <i>et al.</i> (2013) |
| | 505.III.3 | – | <i>Chiloglot s dyphilla</i> | HM196792 | Roche <i>et al.</i> (2010) |
| | SRBG01.II.3 | Australian National Botanical Garden, Acton, ACT, Australia | <i>Chiloglot s trapeziformis</i> | HM196793 | Roche <i>et al.</i> (2010) |
| <i>T. pruinosa</i> | DAOM 17641 | Ontario, Canada | Sporophore on <i>Populus</i> sp. | AY373295 | McCormick <i>et al.</i> (2004) |
| <i>T. punctata</i> | CLM 2012 | Northern Sydney, NSW, Australia | <i>Cryptostylis subulata</i> root | MT008124 | Arif <i>n et al.</i> (2020) |
| | CLM 2017 ^f | Northern Sydney, NSW, Australia | <i>Cryptostylis subulata</i> root | MT008122 | Arif <i>n et al.</i> (2020) |
| | CLM 2018 | Northern Sydney, NSW, Australia | <i>Cryptostylis subulata</i> root | MT008121 | Arif <i>n et al.</i> (2020) |
| <i>T. rosea</i> | CLM 1770 | WA, Australia | <i>Spiculaea ciliata</i> root | MN947568 | Arif <i>n et al.</i> (2020) |
| | CLM 1773 ^f | WA, Australia | <i>Spiculaea ciliata</i> root | MN947569 | Arif <i>n et al.</i> (2020) |
| | CLM 1774 | WA, Australia | <i>Spiculaea ciliata</i> root | MN947570 | Arif <i>n et al.</i> (2020) |
| <i>T. secunda</i> | CLM 274 | Talbot, WA, Australia | <i>Paracaleana triens</i> | KF476580 | Linde <i>et al.</i> (2013) |
| | CLM 222 | Talbot, WA, Australia | <i>Paracaleana minor</i> | KF476568 | Linde <i>et al.</i> (2013) |
| | CLM 009 ^f | Talbot, WA, Australia | <i>Drakaea elast ca</i> | KF476575 | Linde <i>et al.</i> (2013) |
| <i>Tulasnella</i> sp. ECU 5 | DC 225 | Ecuador | Branch | KC152397 | Cruz <i>et al.</i> (2014) |
| | DC 225 | Ecuador | Branch | KC152398 | Cruz <i>et al.</i> (2014) |
| <i>Tulasnella</i> sp. ECU 6 | DC 185 | Ecuador | Branch | KC152401 | Cruz <i>et al.</i> (2014) |

Table 1. (Cont nued).

| Species | Strain no. | Origin | Host | GenBank accession no. | Reference |
|-----------------------|----------------------|--|---|----------------------------|--------------------------------|
| <i>Tulasnella</i> sp. | DC 262 | Ecuador | Branch | KC152409 | Cruz <i>et al.</i> (2014) |
| | DC 294 C009 | Germany | Rot en wood | KC152387 | Cruz <i>et al.</i> (2014) |
| | DC 294 C016 | Germany | Rot en wood | KC152394 | Cruz <i>et al.</i> (2014) |
| <i>T. sphagnet</i> | CLM 084 | Australia | <i>Arthrochilus oreophilus</i> | KF476594 | Linde <i>et al.</i> (2013) |
| | CLM 085 | Australia | <i>Arthrochilus oreophilus</i> | KF476595 | Linde <i>et al.</i> (2013) |
| | 12033.1 [†] | Kosciuszko NP ^o NSW, Australia | <i>Chiloglot saf. valida</i> | KY095117 | Linde <i>et al.</i> (2017) |
| <i>T. tomaculum</i> | 13102.1 | Kosciuszko NP ^o NSW, Australia | <i>Chiloglot s turfosa</i> | KY445924 | Linde <i>et al.</i> (2017) |
| | 13065.2 | Kosciuszko NP ^o NSW, Australia | <i>Chiloglot ssp.</i> | KY445925 | Linde <i>et al.</i> (2017) |
| | KC 429 | – | – | AY373292 | McCormick <i>et al.</i> (2004) |
| <i>T. violea</i> | K(M)123675 | England, UK | – | KC152380 | Cruz <i>et al.</i> (2014) |
| | DC 177 | Ecuador | Decaying wood | KC152414 | Cruz <i>et al.</i> (2014) |
| <i>T. warcupii</i> | DC 292 | Germany | Decaying wood | KC152412 | Cruz <i>et al.</i> (2014) |
| | DC 292 | Germany | Decaying wood | KC152435 | Cruz <i>et al.</i> (2014) |
| | DC 293 | Germany | Decaying wood | KC152437 | Cruz <i>et al.</i> (2014) |
| | KC 851 | – | – | AY373293 | McCormick <i>et al.</i> (2004) |
| | K(M) 164256 | England, UK | Underside of <i>Fagus sylvatica</i> log | KC152411 | Cruz <i>et al.</i> (2014) |
| | CLM 007 | Atherton, Tablelands, QLD, Australia | <i>Arthrochilus oreophilus</i> | KF476600 | Linde <i>et al.</i> (2013) |
| CLM 022 | Australia | <i>Arthrochilus oreophilus</i> | KF476601 | Linde <i>et al.</i> (2013) | |
| CLM 028 | Australia | <i>Arthrochilus oreophilus</i> | KF476599 | Linde <i>et al.</i> (2013) | |

[†]Indicates type specimens.

Table 2. Gene sequences (28S) retrieved from GenBank, and newly generated in this study.

| Species | Strain no. | Origin | Host | GenBank accession no. | Reference |
|---------------------------------|--------------------------|---|------------------------------------|-----------------------|--------------------------------|
| <i>Agaricus bisporus</i> | CBS 151.46 | – | – | MH867670 | Vu <i>et al.</i> (2019) |
| <i>Botryobasidium botryosum</i> | AFTOL-ID 604 | – | – | DQ089013 | Nilsson, unpublished |
| <i>Bo. isabellinum</i> | GEL2109 | – | – | AF393047 | Nilsson, unpublished |
| <i>Bo. subcoronatum</i> | GEL 1286 | – | – | AF287850 | Binder & Hibbet (2002) |
| <i>Burgella lutea</i> | Etayo 27623 [†] | Bolivia | Cort colour lichens | KC336075 | Diederich <i>et al.</i> (2014) |
| <i>Bu. favoparmeliae</i> | JL192-01 | Oklahoma, USA | <i>Flavoparmelia balt morensis</i> | DQ915469 | Lawrey <i>et al.</i> (2007) |
| <i>Burgoa moriformis</i> | VCH 33 [†] | Inisherik, Crom, Fermanagh, Ireland | <i>Salix</i> bark | DQ915477 | Lawrey <i>et al.</i> (2007) |
| <i>Cantharellus addaiensis</i> | BB 96.010 | Zambia | – | KM484680 | Shao <i>et al.</i> (2014) |
| <i>Ca. altipes</i> | BB 07.019 [†] | USA | – | KF294627 | Buyck <i>et al.</i> (2014) |
| <i>Ca. amethysteus</i> | 993/estades | – | – | MG450679 | Buyck <i>et al.</i> (2018) |
| <i>Ca. cibarius</i> | CC15SWE | Sweden | <i>Betula</i> | JX030441 | Foltz <i>et al.</i> (2013) |
| <i>Ca. ferruginasecens</i> | GE sn | France | – | KM484681 | Shao <i>et al.</i> (2014) |
| <i>Ca. formosus</i> | BB 13.163 | USA | – | KM484683 | Shao <i>et al.</i> (2014) |
| <i>Ca. lateritus</i> | JJNC-Canth 2 | USA | – | KM484686 | Shao <i>et al.</i> (2014) |
| <i>Ca. longisporus</i> | ER 107 | Madagascar | – | KM484688 | Shao <i>et al.</i> (2014) |
| <i>Ca. minor</i> | BB 07.057 | USA | – | KF294632 | Buyck <i>et al.</i> (2014) |
| <i>Ca. texensis</i> | BB 07.018 [†] | USA | – | KF294626 | Buyck <i>et al.</i> (2014) |

Table 2. (Cont nued).

| Species | Strain no. | Origin | Host | GenBank accession no. | Reference |
|---------------------------------------|-------------------------|--|------------------------------------|-----------------------|-----------------------------------|
| <i>Ceratobasidium bulbillifaciens</i> | Eichler-Cezanne 8193 | Germany | Bark of <i>Acer platanoides</i> | KC336073 | Diederich <i>et al.</i> (2014) |
| <i>C. bulbillifaciens</i> | Eichler-Cezanne 8067 | – | Bark of <i>Fraxinus</i> | KC336071 | Diederich <i>et al.</i> (2014) |
| <i>C. globisporum</i> | CBS 569.83 ^T | Queensland Australia | – | MH873365 | Vu <i>et al.</i> (2019) |
| <i>C. pseudocornigerum</i> | CBS 568.83 ^T | Australia | – | MH873364 | Vu <i>et al.</i> (2019) |
| <i>C. ramicola</i> | Java 11 | Java | <i>Theobroma cacao</i> | HQ424243 | Samuels <i>et al.</i> (2012) |
| <i>C. theobromae</i> | South Sulawesi 1 | South Sulawesi | Pet ole of <i>Theobroma cacao</i> | KU319575 | Samuels <i>et al.</i> (2012) |
| | South Sulawesi 6 | South Sulawesi | Pet ole of <i>Theobroma cacao</i> | KU319577 | Samuels <i>et al.</i> (2012) |
| | South Sulawesi 11 | South Sulawesi | <i>Theobroma cacao</i> | HQ424241 | Samuels <i>et al.</i> (2012) |
| | South Sulawesi 10 | South Sulawesi | <i>Theobroma cacao</i> | HQ424242 | Samuels <i>et al.</i> (2012) |
| <i>Ceratorhiza oryzae-sat vae</i> | CBS 439.80 | Japan | <i>Oryza sat va</i> | MH873047 | Vu <i>et al.</i> (2019) |
| <i>Clavulina amazonensis</i> | AMV1973 | Colombia | <i>Pseudomonotes tropenbosii</i> | KT724123 | Vasco-Palacios (2016) |
| <i>Cl. cinerea</i> | KHL 11694 | Lammi, Finland | – | AM259211 | Nilson <i>et al.</i> (2006) |
| <i>Cl. cf. cristata</i> | BB 12.083 | Italy | – | KM484694 | Shao <i>et al.</i> (2014) |
| <i>Cl. purpurascens</i> | ZP-3065 | China | Soil | MK564124 | Wu <i>et al.</i> (2019) |
| <i>Clavucilium delectabile</i> | KHL 11147 | Norway | – | AY586688 | Larsson <i>et al.</i> (2004) |
| <i>Craterellus cinereof mbratus</i> | JOH4 | Columbia | <i>Pseudomonotes tropenbosii</i> | KT724159 | Vasco-palacios (2016) |
| <i>Cr. lutescens</i> | BB 13.048 | Canada | – | KM484696 | Shao <i>et al.</i> (2014) |
| <i>Cr. tubaeformis</i> | BB 1324 | USA | – | KM484697 | Shao <i>et al.</i> (2014) |
| | BB 07.293 | Slovakia | – | KF294640 | Buyck <i>et al.</i> (2014) |
| <i>Dacrymyces chrysospermus</i> | FPL11353 | – | – | AF287855 | Hibbet <i>et al.</i> (2000) |
| <i>D. st llatus</i> | CBS 195.48 | France | – | MH867857 | Vu <i>et al.</i> (2019) |
| <i>Gloeotulasnella cyst diophora</i> | KW 2871 | – | – | AY585831 | Shef erson <i>et al.</i> (2005) |
| <i>Haplotrichum conspersum</i> | AFTOL ID 1766 | – | – | DQ521414 | Matheny <i>et al.</i> unpublished |
| <i>Hydnum elatum</i> | FRI62309 | Kampung Jelawat-Tasik Bera Pahang Malaysia | – | KU612691 | Feng <i>et al.</i> (2016) |
| <i>H. elliposporum</i> | FD3281 | Switzerland | – | KX086217 | Beenken, unpublished |
| <i>H. magnorufescens</i> | 161209 | Slovenia | – | KU612669 | Feng <i>et al.</i> (2106) |
| <i>H. rufescens</i> | BB 07.340 | Slovakia | – | KM484698 | Shao <i>et al.</i> (2014) |
| <i>H. versterholt i</i> | HKAS92342 | Yulong snow mountain, Yunnan, China | – | KU612646 | Feng <i>et al.</i> (2016) |
| <i>Minimedusa obcoronata</i> | CBS 120605 | Thailand | <i>Eucalyptus camaldulensis</i> | GQ303309 | Cheewangkoon <i>et al.</i> (2009) |
| <i>Mult davula mucida</i> | DSH96-056 | – | – | AF287875 | Hibbet <i>et al.</i> (2000) |
| <i>M. mucida</i> | TUB 011734 | – | – | EU909345 | Krause <i>et al.</i> (2011) |
| <i>M. vernalis</i> | GB-BN-1 | Sweden | – | AM259214 | Nilsson <i>et al.</i> (2006) |
| <i>Rhizoctonia f occosa</i> | CBS 337.36 ^T | Indonesia | – | MH867319 | Vu <i>et al.</i> (2019) |
| <i>R. quercus</i> | CBS 313.35 ^T | Italy | Root of <i>quercus pedunculata</i> | MH867202 | Vu <i>et al.</i> (2019) |
| <i>R. repens</i> | CBS 298.32 | Netherlands | <i>Orchis morio</i> | MH866781 | Vu <i>et al.</i> (2019) |
| <i>Schizophyllum commune</i> | MUT 4875 | Mediterranean Sea | <i>Flabellia pet olata</i> | MF115832 | Poli <i>et al.</i> (2018) |
| <i>Sistotrema adnatum</i> | FCUG 700 | – | – | DQ898699 | Moncalvo <i>et al.</i> (2006) |
| <i>S. biggisae</i> | FCUG 782 | – | – | DQ898697 | Moncalvo <i>et al.</i> (2006) |
| <i>S. coronilla</i> | AFTOL-ID 618 | – | – | DQ457641 | Moncalvo <i>et al.</i> (2006) |

Table 2. (Cont nued).

| Species | Strain no. | Origin | Host | GenBank accession no. | Reference |
|---------------------------------|----------------------------|-------------------------------------|--|-----------------------|-------------------------------|
| <i>S. hypogaeum</i> | CBS 394.63 ^T | Australia | Soil | MH869926 | Vu et al. (2019) |
| <i>S. oblongisporum</i> | FCUG 2117 | – | – | DQ898703 | Moncalvo et al. (2006) |
| <i>S. sernanderi</i> | CBS 926.70 | – | – | AF518650 | Hibbet & Binder (2002) |
| <i>Sistotremella brinkmanii</i> | CBS 186.39 | Michigan, USA | – | MH867474 | Vu et al. (2019) |
| <i>Sistotremella perpusilla</i> | CBS 126048 | North Carolina, USA | <i>Abies</i> | MH875516 | Vu et al. (2019) |
| <i>Thanatephorus cucumeris</i> | AFTOL-ID 2022 | – | – | DQ917658 | Matheny et al. unpubl. |
| | CBS 340.51 | England, UK | – | MH868410 | Vu et al. (2019) |
| <i>Tremella macrobasidiata</i> | AM453 | Portugal | <i>Lecanora chlorotera</i> | KT334595 | Zamora et al. (2016) |
| <i>Tr. mesenterica</i> | AM30 | Wedin, Sweden | – | JN043569 | Millanes et al. (2011) |
| | CBS:6973 ^T | Vancouver, British Columbia, Canada | <i>Alnus rubra</i> | KY109900 | Vu et al. (2016) |
| <i>T. anat cula</i> | UAMH 5428 | Alberta, Canada | Roots of <i>Calypso bulbosa</i> | AY243520 | Taylor et al. (2003) |
| <i>T. asymmetrica</i> | MAFF P305806 | | <i>Thelymitra luteocilium</i> | DQ388046 | Suarez et al. (2006) |
| <i>T. aurant aca</i> | DAOMC 251988 | Pennsylvania, USA | Rot en wood | MK627511 | This study |
| | DAOM 970795 | | | | |
| | DAOMC 251989 | Tennessee, USA | Rot en wood and <i>Crepidotus</i> spp. | MK627512 | This study |
| | PBM 4158 | | | | |
| | DAOMC 252083 | Victoriaville, Quebec, Canada | <i>Fomitopsis betulina</i> | MK627513 | This study |
| | DAOM 970821 | | | | |
| | DAOMC 252084 | Ot awa, Ontario, Canada | Rot en <i>Populus</i> wood | MK627514 | This study |
| | DAOM 970822 | | | | |
| | DAOMC 252086 | Montreal, Quebec, Canada | Rot en wood | MK627515 | This study |
| | DAOM 970819 | | | | |
| | DAOMC 252085 | Ot awa, Ontario, Canada | Rot en wood | MK627516 | This study |
| | DAOM 970820 | | | | |
| <i>T. calospora</i> | SPRR.R2 | India | <i>Paphiopedilum druryi</i> | MN271391 | Parthibhan & Rammasubu (2020) |
| <i>T. eremophila</i> | 13062 MD | – | <i>Euphorbia of cinarum</i> | KJ701189 | Crous et al. (2015) |
| <i>T. irregularis</i> | CBS 574.83 | NT, Australia | <i>Dendrobium dicuphum</i> | AY243519 | Taylor et al. (2003) |
| <i>T. phuhinrongklaensis</i> | SDBR-CMU-CR41 ^T | – | – | MF427703 | Rachanarin et al. (2018) |
| | SDBR-CMU-CR42 | – | – | MF427704 | Rachanarin et al. (2018) |
| | SDBR-CMU-CR43 | – | – | MF427705 | Rachanarin et al. (2018) |
| | SDBR-CMU-CR44 | – | – | MF427706 | Rachanarin et al. (2018) |
| <i>T. pruinosa</i> | DAOM 17641 | Richmond Hill, Ontario, Canada | <i>Populus</i> sp. | AF518662 | Hibbet & Binder (2002) |
| <i>Tulasnella</i> sp. | GEL 4461 | – | – | AJ406436 | Langer (2001) |
| | GEL 4745 | – | – | AJ406436 | Langer (2001) |
| | GEL 5130 | – | – | DQ898731 | Moncalvo et al. (2006) |
| <i>T. violea</i> | DAOM 222001 | – | – | AY293216 | Binder et al. (2005) |
| | AFTOL-ID 1879 | – | – | DQ520097 | Garnica & Weiß, unpublished |
| <i>Ust lago maydis</i> | CBS 358.32 | – | – | MH866814 | Vu et al. (2019) |

^TIndicates type specimens or ex-type strains.

Table 3. Primer names and sequences used for this study, including the new primers ITS5_hormo, ITS4_hormo and LROR_hormo and the *Tulasnella*-specific ITS4 primer, ITS4_tul, with melting temperature (T_m) in degrees Celsius. Point mutations are in bold, nucleotide additions are in bold and underlined and nucleotide deletions are shown as a blank space with a bold underline.

| | Sequence (5' to 3') | T_m (°C) | Reference |
|-------------------|------------------------|------------|---------------------------------------|
| V9G | TTACGTCCTGCCCTTTGTA | 56 | de Hoog & Gerrits van den Ende (1998) |
| ITS5 | GGAAGTAAAAGTCGTAACAAGG | 51 | White <i>et al.</i> (1991) |
| ITS5_hormo | GGAAGTACAAGTCGTAACAAGG | 53 | This study |
| ITS4 | TCCTCCGCTTATTGATATGC | 52 | White <i>et al.</i> (1991) |
| ITS4_tul | CCGCCAGATTCACACATTGA | 55 | Taylor & McCormick (2008) |
| ITS4_hormo | TCCTCCGCTGAATAATATGC | 52.1 | This study |
| LROR | ACCCGCTGAACTTAAGC | 52 | Moncalvo <i>et al.</i> (2000) |
| LROR_hormo | ACCCGCTIGA_TTTAAGC | 50 | This study |
| LR3 | CCGTGTTTCAAGACGGG | 53 | Moncalvo <i>et al.</i> (2000) |
| LR3R | GTCTGAAACACGGACC | 50 | Moncalvo <i>et al.</i> (2000) |
| LR5 | TCCTGAGGGAACTTCG | 51 | Moncalvo <i>et al.</i> (2000) |
| LR8 | CACCTTGGAGACCTGCT | 54 | Hopple & Vilgalys (1999) |

Our initial attempts to amplify ITS and 28S separately from cultures of *H. aurantaca* using universal primers, or a putative *Tulasnella*-specific ITS primer (Taylor & McCormick 2008), were unsuccessful, with only faint or multiple bands visualized. Diverse techniques such as annealing point gradients from 60–55 °C, a touchdown PCR of 40 cycles each at 57 and 55 °C, and altering primer and DNA concentrations were all unsuccessful in amplifying the DNA. This was solved using the forward primer V9G, slightly upstream from the ITS locus at the end of the 18S (de Hoog & Gerrits van den Ende 1998) along with the reverse primer LR3, slightly downstream from the LROR primer on the 28S (Raja *et al.* 2017). Sequences of this fragment showed that *H. aurantacus* DAOMC 251989 has mismatched binding sites in the ITS4, ITS5, ITS1F and LROR priming regions. For subsequent amplification and sequencing, we designed the new primers ITS4_hormo, ITS5_hormo and LROR_hormo (Table 3), correcting the mismatches identified by our first sequence. This enabled the routine amplification and sequencing of both markers.

The sequences of our six strains of *Hormomyces* were very similar. For the ITS (Fig. 2), four of the strains had identical sequences; DAOMC 251989 had a single nucleotide insertion, and DAOMC 252084 differed from the other strains by 3–4 nucleotides. For the 28S (Fig. 3), the differences between each strain were no more than two nucleotides, resulting in similarities > 99.5%.

All analyses of both genes confirmed the phylogenetic relationship with *Tulasnellaceae*, *Cantharellales*, suggested by the initial 18S sequence, remote from the previously proposed relatives of *H. aurantacus* in *Tremellales* or *Dacrymycetales*. This directed our sampling of taxa to test whether *Hormomyces* was nested within or distinct from existing genera of *Tulasnellaceae*. All BI and ML analyses resulted in all strains of *H. aurantacus* forming a monophyletic clade nested within *Tulasnella* and separate from other genera of *Cantharellales*. In the ITS analysis, *H. aurantacus* belongs with high support to a clade including various *Tulasnella* species including *T. violea*, *T. eichleriana* and several orchid root endophytes. Unfortunately our BI analysis of the ITS loci tends to result in polytomy for this clade, although each species remains phylogenetically coherent and distinct. The exact sister group of *T. aurantaca* in this clade is unclear. A phylogenetic analysis of the ITS locus by Arifin *et al.* (2020),

which used sequences of *H. aurantacus* obtained during our study, also suggested the placement of *H. aurantaca* in the same clade within *Tulasnella*, but sister to two unidentified *Tulasnella* species from Ecuador.

Our 28S analysis also nests *H. aurantacus* within *Tulasnella*, suggesting it is sister to *Tulasnella* sp. GEL4461, with a difference of 47 bp (94.6% similarity). Based on a MegaBLAST search of GenBank, the closest identified match for the LSU locus was *Tulasnella obscura* (AJ406435, identities = 485/517 (94%), 7 gaps (1%) with a 57% query cover). Other matches were less than 85% similar and with an E value below 1e-100.

Based on our analyses, *Tulasnella* appears to be monophyletic (with the exception of *Tulasnella eremophila*), forming two distinct clades with high statistical support in the ITS phylogeny, and possibly four distinct clades in the 28S phylogeny. Because only six species of *Tulasnellaceae*, including our own, have sequences for both ITS and 28S, a concatenated analysis was not attempted. All of the asexual species previously referred as *Epulorhiza* currently sequenced using the ITS and LSU loci are easily separated from *H. aurantacus*, suggesting that *H. aurantacus* is a distinct species of *Tulasnella*.

Comparison of herbarium specimens of *H. aurantiacus*, *H. fragiformis* and *H. callorioides*

All of the examined specimens identified as *H. aurantacus*, *H. fragiformis* or *H. callorioides* had similar micromorphology and no distinct separation was detected based on an extensive analysis of conidial sizes (Supplementary Fig. S1). Measurements overlapped with the lectotype specimen of *H. fragiformis* [(7–)7.5–9.5(–11) × 6–7.5(–8.5) μm (mean 8.66 ± 0.1 × 6.75 ± 0.1, Q 1.29, n = 50)] and the holotype of *H. callorioides* [(7–)7.5–9.5(–11) × (5–)5.5–6.5(–7) μm (mean 8.45 ± 0.1 × 6.21 ± 0.1, Q 1.37, n = 50)], with the average conidial size for all examined specimens being 8.43 ± 0.1 × 6.37 μm ± 0.1 (Supplementary Fig. S2).

Although the protologue of *H. callorioides* described its sporodochia as pink (Kalchbrenner & Cooke 1880), our examination of the holotype showed dried sporodochia that were terracotta to dark brown. The original pigments may have degraded over the years or darkened with preservation. The colours observed on the lectotype of *H. fragiformis* designed by McNabb (1969)




Fig. 2. Phylogenetic tree for species of *Tulasnella* and *Ceratobasidium* for the ITS region, based on Bayesian inference analysis using MrBayes. The values above the branches are Bayesian posterior probabilities/maximum likelihood bootstrap values. Bootstrap values of 70 % and Bayesian posterior probabilities of 0.70 are shown with bootstrap values of 100 % and Bayesian posterior probability of 1.00 replaced by an asterisk (*). ^T indicates type specimens.

and holotype of *H. callorioides*, both in dry specimens and after rehydration, were within the same range as that we saw in our own specimens and those preserved in BPI and DAOM. We found no evidence to support a hypothesis that sporodochial colour can be used to distinguish these three putative species when the specimens were examined side by side. Similarly, there were no other morphological or microscopic characters that allowed the distinction of the several specimens identified as *H. fragiformis* or *H. aurantacus* or the holotype of *H. callorioides*, despite the disjunct geographical locations among the three species.

Taxonomy

The genus *Tulasnella* was proposed for protection against *Epulorhiza* and *Hormomyces* by Stalpers et al. (2021), in accordance of the Shenzhen code (Turland et al. 2018). The remaining names in *Epulorhiza* were transferred to *Tulasnella* there. Here, the generic description of *Tulasnella* is emended to include the asexual morph characters for taxa described in *Hormomyces* and *Epulorhiza*.

Tulasnella J. Schröt., *Kryptogamen-Flora von Schlesien* 3.1(25–32): 397. 1888. **nom. cons. prop.**

- Synonyms:* *Hormomyces* Bonord., *Handb. Allgem. mykol.* (Stuttgart): 150. 1851 (asexual synonym).
- Prototremella* Pat., *J. Bot., Paris* 2: 269. 1888, *f de Donk* 1966.
- Pachysterigma* Johan-Olsen ex Bref., *Unters. Gesamtgeb. Mykol.* (Liepzig) 8: 5. 1888, *f de Donk* 1966.
- Muciporus* Juel, *Bih. K. svenska VetenskAkad. Handl., Afd.*: 23. 1897, *f de Donk* 1966.
- Gloeotulasnella* Höhn. & Litsch., *Wiesner Festschrif* (Wien): 57. 1908, *f de Donk* 1966.
- Hormisciopsis* Sumst., *Mycologia* 6: 32. 1914 (asexual synonym).
- Epulorhiza* R.T. Moore, *Mycotaxon* 29: 94. 1987 (asexual synonym).

Typifcat on: *Tulasnella lilacina* J. Schröt. 1888.

Basidiomes (when present) crust-like, of en on rotten wood, leaves or litter, of en pinkish or purple. *Hymenium* composed




Fig. 3. Phylogenetic tree for species of *Cantharellales* including species of *Tulasnella* for the 28S gene, based on Bayesian inference analysis using MrBayes. The values above the branches are Bayesian posterior probabilities/maximum likelihood bootstrap values. Bootstrap values of 70% and Bayesian posterior probabilities of 0.70 are shown with bootstrap values of 100% and Bayesian posterior probability of 1.00 replaced by an asterisk (*). † indicates type specimens.

of hyphae with or without clamps depending on the species, subhymenial structures of an absent. *Basidia* with four sterigmata that are strongly swollen at the base, each separated by a septum from the clavate basal cell, which often collapses. *Basidiospores* variably shaped, from globose to helicoid; secondary conidia occasionally produced on germinating primary spores (Ingold 1984). Septa with central dolipores with continuous parentheses. *Asexual morphs* (when present) forming mycorrhizae with orchid roots or liverworts, or conspicuous orange to red pustulate, gelatinous, sporodochia on rotten wood. *Stroma* absent. *Conidia* in blastoacropetal chains, which are dichotomously branched in some species, and arise directly from submerged hyphae or in sporodochia. Conidia aseptate, smooth or with minute pits, globose, subglobose, ellipsoidal to barrel-shaped, aseptate, hyaline, sometimes with orange droplets. Somatic hyphae dikaryotic (Currah *et al.* 1990), lacking clamp connections when associated with asexual structures. Sclerotia occasionally produced.

Notes: The sexual morphs of *Tulasnella* are well-characterized by the production of thin, resupinate basidiomes on decaying substrates. They have unique "tulasnelloid" basidia with a basally swollen sterigma separated by a septum from the club-shaped basal cell, which usually collapses (Roberts 1994). Ultrastructure of dolipore septa is often used as a character to support class-level classification of *Basidiomycota* (Celio *et al.* 2006). Comparatively few exemplars are studied for each class and the character is rarely used for genus or species level classifications, which tend to be supported by DNA sequencing (*e.g.* Almeida *et al.* 2014, Linde *et al.* 2017, Arifin *et al.* 2020). We did not examine septal ultrastructure in this study. Most known asexual morphs of *Tulasnella* are characterized by the production of acropetal chains of conidia, and for most of the known asexual species, a relationship with orchids. *Tulasnella aurantiaca* is the exception because it grows on wood.

Apart from the species formerly included in *Epulorhiza*, a few other asexual states are described for other species of *Tulasnellaceae* but lack the characteristic monilioid conidial

chains of *T. aurantaca*. *Tulasnella valentini*, which grows on rotten wood, is reported to produce single apiculate, obclavate or fusiform conidia terminating irregularly lanceolate conidiogenous cells (Van de Put & Antonissen 1996). This is an unusual character state for a member of *Tulasnellaceae* and this asexual-sexual connection needs to be confirmed experimentally to eliminate the possibility that the observed conidia might have been those of a mycoparasite; for that reason, we did not include these deviating character states in the generic diagnosis above. *Stilbotulasnella conidiophora*, described from palm fronds but to date unsequenced, can be distinguished by its synnematosous conidiomata and ellipsoidal asexual conidia produced in slimy masses from percurrently proliferating conidiogenous cells (Bandoni & Oberwinkler 1982).

Seifert et al. (2011) synonymized the monotypic *Hormisciopsis* with *Hormomyces* based on the protologue, and we include it as a synonym of *Tulasnella* above. We did not examine specimens of *Hormisciopsis gelatinosa*, but the protologue (Sumstine 1914) provides observations that are identical to what we have seen in specimens of *T. aurantaca*. Seifert et al. (2011) followed von Höhnelt (1917) and Donk (1962) in synonymizing *Sphaerocolla* with *Hormomyces*. However, re-examination of the protologue and slides from the holotype of *S. aurantaca* (on *Betula*, Mustala, June; H herb. Karsten PAK 3341; Supplementary Fig. S3), the type species of *Sphaerocolla*, suggests this was an error. Although the micromorphology is similar, the conidia of *S. aurantaca* are slightly oblate rather than slightly ellipsoidal, and the fungus is reminiscent of the poorly-documented yeast *O. margariferum*, which occurs on slime fluxes on trees (Kurtzman 2011), a similar ecological niche to that reported for *S. aurantaca* on living *Betula* trees. A notable difference is that *O. margariferum* produces endospores, which we did not observe in the material of *S. aurantaca*, but these are apparently produced only in culture (Smith 1997) and no cultures of *S. aurantaca* were isolated. Therefore, *S. aurantaca* should be considered distinct from *T. aurantaca* and may be conspecific with *O. margariferum*.

Based on our nuclear DNA phylogenies, *Hormomyces* is congeneric with *Tulasnella*, but appears to be distinct from all sequenced species of the latter genus. Therefore, a new combination is proposed:

Tulasnella aurantiaca (Bonord.) J. Mack & Seifert, **comb. nov.**
Mycobank MB832426. Fig. 4.

Synonyms: *Hormomyces aurantacus* Bonord., *Handb. Allgem. mykol.* (Stuttgart): 150. 1851.

Typification: fig. 234, taf. XI, in Bonorden 1851, *Handb. Allgem. mykol.* (Stuttgart): (lectotype, proposed here MBT388543, reproduced here as Fig. 1).

Hypsilophora callorioides Kalchbr. & Cooke, *Grevillea* 9: 18. 1880.
Hormomyces callorioides (Kalchbr. & Cooke) Sacc., *Syll. fung.* (Abellini) 6: 813. 1888.

Hypsilophora fragiformis Cooke, in Farlow, *Appalachia* 3: 247. 1884

Hormomyces fragiformis (Cooke) Sacc., *Syll. fung.* (Abellini) 6: 182. 1888.

Hormisciopsis gelatinosa Sumst., *Mycologia* 6: 32. 1914.

Basidiomes unknown. **Sporodochia** effuse, often pustulate, confluent in masses up to 5 cm long, occasionally more, or rarely solitary and <1 cm long, gelatinous or cartilaginous when fresh,

waxy when dried, colours variable, from deep orange to garnet red when fresh, ranging from blonde to brown with various shades of orange and red when dry. **Hyphae** immersed, septate, branched, 2–4 µm wide, dikaryotic, binucleate (Supplementary Fig. S4), clamp connections absent, no stroma formed. **Conidiophores** arising from hyphal cells, clamp connection absent. **Conidia** blastoc, in moniloid, branched acropetal chains, branching bifurcate, pattern variable, occurring mainly near the base of the chains, with usually fewer than five conidia between bifurcations, the terminal chains typically longer, often with up to 15 conidia and occasionally more, chains not readily separating into individual conidia. Conidia hyaline, often with conspicuous orange guttules when fresh, aseptate, smooth, subglobose to ellipsoidal or occasionally globose, often truncate, variable in length and width, (4.5–)7.5–9.5(–13) × (4–)5.5–7(–8.5) µm (mean 8.4 ± 0.1 × 6.4 ± 0.1, Q 1.31), thick-walled with walls ~1 µm thick.

Colony diameter after 14 d, on MEA with near-UV (spectral range 300–400 nm) 30–50 mm on MEA in darkness 30–50 mm on CMA with near-UV 25–45 mm, on CMA in darkness 30–50 mm, on OA with near-UV 55–65 mm, on OA in darkness 40–55 mm. **Colonies** flat, often immersed, filamentous, often circular or irregular with undulate margins. **Hyphae** 2–5 µm wide. **Sporodochia** produced after 1–2 wk abundantly on OA, and in some strains on MEA, sterile on CMA, concolorous with mycelium, or more vibrantly coloured, especially when exposed to light: on CMA with near-UV after one month Salmon (6A4) to Pinkish White (9A2) in four strains and Cognac (6E7) in one; on CMA in darkness white, Blonde (5C4), Pinkish White (9A2) or Cognac (6E7); on MEA with near-UV white in four strains and Tangerine (6B7) in one; on MEA in darkness white in four strains and Yellowish White (4A2) in one; on OA under near-UV Greyish Red (7B4) to Pastel Red (8–9A5); on OA in darkness Pale Yellow (3A3) to Orange White (5A2) or Pale Orange (6A3). Conidia similar to those *in vivo* but slightly longer and narrower, (6.5–)7.5–10(–12.5) × (4–)5–6.5(–8) µm (mean 9 ± 0.5 × 5.8 ± 0.3, Q 1.57 ± 0.1).

Cardinal temperatures: Optimum 25 °C, minimum <5 °C, maximum 30–37 °C. Growth does not resume in cultures left for 1 mo at 37 °C, when moved to an incubator at 20 °C.

Distribution: Widely distributed in eastern North America, known from Ontario and Québec south to North Carolina and Tennessee and westward to Ohio. *Tulasnella aurantaca* is known from western North America (one specimen collected in Arizona), Europe (Austria, Germany) and South Africa (Western Cape).

Habitat: Lignicolous and apparently saprobic, reported on rotten wood of conifers (*Thuja*) and angiosperms such as species of *Populus*, *Platanus*, *Liquidambar* and *Vaccinium*. It sometimes overgrows other fungi such as *Fomitopsis betulina* and *Crepidotus* spp.

Specimens examined (*indicates specimens that were cultured): As *Hormomyces aurantacus*: Austria, Salzburg, Salzburg, on bark of rotten wood, 1880, E.A. Rau (BPI 726623). Canada, Ontario, Ottawa, Portobello Park, on rotten wood, probably *Populus* sp., Aug. 2014, J. Mack (DAOM 970822); *ibid.*, Jul. 2015 (*DAOMC 252084); Ottawa, on rotten wood, probably *Populus* sp., 15 Jul. 2017, J. Mack (DAOM 970820, *DAOMC 252085); Quebec, Montreal, on rotten wood, 25 Oct. 2018, G. Carter, isol. J. Mack (DAOM 970818, *DAOMC 252086); Victoriaville, on *Piptoporus*




Fig. 4. *Tulasnella aurantaca*. A, B. Appearance of fresh sporodochia on rot en wood. C. Detail of dried sporodochia. D. Detail of rehydrated sporodochia. E–G: 28-d-old culture, E on CMA with near-UV, F on OA with near-UV, G on OA in darkness. H, K. Living conidial chains in water. I, J. Conidial chains from culture on OA for 27 d under near-UV. L. Conidial chains in lact c acid. Scale bars: C–D = 500 µm, H, K–L = 10 µm, I, J = 20 µm.

betulinus, 13 Jun. 2016, *B. Fort* et al. isol. *J. Mack* (DAOM 970821, *DAOMC 252083). **South Africa**, on bark of unidentified tree, undated, *V. Duthie Augusta* (BPI 702911). **USA**, Arizona: Portal, Greenhouse Canyon, on rotten wood of *Platanus* sp., 26 Aug. 1956, *J.L. Lowe & R.L. Gilbertson* (BPI 726600); Kentucky, Crittenden, on rotten wood, 11 Jul. 1910, *C.G. Lloyd* (BPI 702908); Maryland, Great Falls, on rotten wood, 10 Oct. 1936, *J.A. Stevenson* (BPI 726621); Great Falls, on rotten log, 17 Oct. 1936, *J.A. Stevenson* (BPI 726619); Patuxent Wildlife Refuge, on wood of *Liquidambar styraciflua*, Jul. 1952, *F. Berry* (BPI 726599); Sligo, on rotten wood and polypores, 16 Oct. 1936, *V.K. Charles & E.E. Dick* (BPI 726619); New York, Adirondack Mountains, Paul Smiths, on rotten wood, 21 Sep. 1927, *H.M. Fitzpatrick* (BPI 702905); Greenport, on bark of *Vaccinium* sp., 3 May 1921, *L. Roy* (BPI 702904); North Carolina, Macon County, Highland, Mirror Lake Rd, on a fallen branch, 27 Aug. 1989, *R.J. Bandoni* no. 8495 (DAOM 970797, as *Hormomyces* sp.). Pennsylvania, Meadville, on rotten wood, 1922, *E.C. Smith* (BPI 702909); no location, on rotten wood, 2018, *D. Newman*, isol. *J. Mack* (*DAOMC 251988); Tennessee, Great Smoky Mountains National Park, Cosby, on rotten wood and *Crepidotus* spp., 27 May 2018, *B.P. Matheny*, isol. *B.P. Matheny* (received as a culture, *DAOMC 251989); West Virginia, Fairmont, on rotten wood, no date, *A. Boulton* (BPI 702903). As *Hormomyces fragiformis*: **Canada**, Ontario, Coopers Falls, on rotten log, 16 Sep. 1952, *R.F. Cain* (DAOM 82339); South of Potageville, on hardwood plank, 6 Jul. 1954, *R.F. Cain* (DAOM 52048); New Durham, on decaying log, 11 Nov. 1930, *R.F. Cain* (DAOM 81729). **USA**, New Hampshire, Shelburne, on dead wood and polypore, Jun. 1883, *M.C. Cooke* K(M) 257483 (holotype); Maryland, Grand Falls, on rotten wood, 11 Oct. 1936, *J.A. Stevenson* (BPI 726622); Missouri, Perryville, Jun. 1883, *C.H. Demetrio* (BPI 726620); North Carolina: Asheville, on rotten wood, 1918, *C.G. Lloyd* (BPI 726625), Winston-Salem, on bark of rotten wood, 4 Jul. 1936, *P.O. Schallert* (BPI 726626); Ohio, Cincinnati, on bark, 3 Oct. 1920, *C.G. Lloyd* (BPI 702914); Chapel Hill, 14 Jan. 1924, on rotting deciduous wood, *J.N. Couch* (UNC 7236, in DAOM); Vermont, Middlebury, on dead wood, 20 Aug. 1896, *E.A. Burt* (BPI 702913); Virginia, W slope of Mt. Elliot, Augusta Co., on rotten wood, 17–21 Jul. 1936, *J.A. Stevenson* (BPI 726617); Rapidan River, Shenandoah National Park, rotten wood, 24 Sep. 1936, *J.A. Stevenson* (BPI 726616); West Virginia, Fairmont, on bark of rotten wood, no date, *A. Boulton* (BPI 626624). As *Hormomyces callorioides*: **South Africa**, Somerset West, on rotting wood, no date, *MacOwan* (holotype K(M) 257481).

Notes: *Tulasnella aurantaca* is characterized by orange to red sporodochia growing on wood or rarely, on other fungi, and the production of ellipsoidal conidia generally shorter than 10 µm. No sexual morph is known. While most species of *Tulasnella* occur on rotten wood (Roberts 1994), until now their asexual morphs have only been reported as symbionts of orchids or liverworts. Most of these symbiotic *Tulasnella* spp. have conidia > 10 µm long. The conidia of *T. aurantaca* are most similar to those of *T. epiphytica*, which are also < 10 µm long and occur in branched, monilioid chains; those of *T. epiphytica*, however, have pitted walls (Pereira et al. 2003). Like *T. aurantaca*, *T. calendulina* also produces orange colonies in culture, but its monilioid cells are larger (Zelmer & Currah 1995). Monilioid chains have been reported from basidiomes of *T. violea*, but the component cells are also >10 µm long (Roberts 1994). Sporodochial asexual morphs resembling *T. aurantaca* include *Oosporidium margariferum* (*Eurotomyces*, *Ascomycota*), which also produces gelatinous sporodochia with long, branched chains of globose conidia, but differs by yeast-like growth in culture and its habit on living woody plants (Kurtzman 2011). *Calloria fusarioides* (*Dermateaceae*, *Helotiales*, *Ascomycota*; asexual morph formerly known as *Cylindrocolla urticae*), also

produces gelatinous, orange sporodochia with long, branched acropetal chains but its conidia are cylindrical (Seifert et al. 2011). *Heteromycephaga glandulosa* (tentatively *Tremellomyces*, cf. Weiß et al. 2014), a parasite of basidiomes of *Exidia glandulosa*, produces clavate conidia attached to conidiogenous cells by a clamp connection (Roberts 1997); it is unclear from the protologue whether the conidia are single or formed in chains.

Tulasnella aurantaca grows easily on standard mycological media and grows *in vitro* between 5 and 30 °C, suggesting that it may be well-adapted to grow in different biomes. Most of the specimens examined were collected between May to October, suggesting that *T. aurantaca* has a long sporulating season, which generally correlates with its optimum growth temperature of 25 °C in temperate climates. However, this species can also occur during the winter, as an examined specimen was collected in January.

After studying the available types of all described *Hormomyces* species and many other specimens, only *H. fragiformis* and *H. callorioides* were appropriately placed in Tubaki's (1976) concept of *Hormomyces*. After careful studies of sporodochial coloration and conidial dimensions (Supplementary Fig. S1), we tentatively consider these two taxa to be conspecific with *H. aurantacus* in agreement with Lloyd (1916). As far as we have been able to determine by microscopy, the herbarium specimens from Europe and South Africa are identical to the North America material. The colour variations we observed by growing cultures on different media in darkness or light, re-vegetation of sporodochia on ca. 30 herbarium specimens, and evaluation of approximately 35–40 photographs of both dried and fresh specimens on Mushroom Observer (Wilson et al. 2020), suggests that the colour distinctions used by previous authors, who examined only a few specimens, may not be diagnostic. However, the taxonomic significance of colour differences should be re-evaluated if additional evidence suggests the existence of cryptic, phylogenetic species within *T. aurantaca*.

The cultures we examined and sequenced were all from North America and form a phylogenetically coherent clade. Bonorden's *H. aurantacus* was described from Europe. Unfortunately, despite the many collecting excursions by the senior author in the Netherlands in the mid-1980s, two visits to South Africa in 1996 and 2006, and requests to both professional and amateur colleagues to watch for this fungus over the past five years, we were unable to obtain fresh specimens or cultures of *T. aurantaca*-like asexual morphs from Europe or Africa. Whether phylogenetically distinct species might occur in other parts of the world remains unknown, and we chose to exercise caution and not to propose any of the currently available material to epitypify *T. aurantaca*. In any case, Bonorden's illustration (reproduced here as Fig. 1), must serve as the lectotype (Art. 9.12, Turland et al. 2018) as proposed in the nomenclator above. If future studies of cryptic species support geographic separation, then *H. fragiformis* may be the appropriate name for material from North America, and *H. callorioides* for specimens from South Africa.

As reviewed in the Introduction, *T. aurantaca* of course was considered the asexual morph of *Tremella mesenterica* (McNabb 1969). Based on our phylogenetic results, the description of the asexual morph of *Tr. mesenterica* by Pipolla & Kotteranta (2008) and the yeast-like rather than filamentous growth of the latter in cultures (Fenwick 1995), this speculative connection is clearly untrue. The sexual morph of *T. aurantaca*, if extant, remains unknown, but would be expected to have typical *Tulasnella* basidiomes and basidia.

Excluded species

Only *H. aurantacus*, *H. fragiformis* and *H. callorioides*, discussed above, conform with the Tubaki (1976) concept of *Hormomyces*, now merged with *Tulasnella*. The remaining described species are discussed here.

Hormomyces abietinus P. Karst., Hedwigia 29: 271. 1890.

Synonym: ? *Dacrymyces deliquescens* Nees, 1816, f de Kennedy 1958.

We did not examine the type specimen. Kennedy (1958) considered *H. abietinus* a synonym of *Dacrymyces deliquescens*. McNabb (1973) synonymized *D. deliquescens* with *D. stllatus* and described the asexual morph of *D. stllatus*, which produces gelatinous, orange sporodochia with chains of arthroconidia $8.16 \times 2.5-5.5 \mu\text{m}$, similar to the conidial dimensions of *H. abietinus* reported by Karsten (1890). Neither Kennedy (1958) nor McNabb (1973) examined the holotype of *H. abietinus* and our request to examine the specimen in Herbarium remained unanswered; the synonymy remains tentative for this reason.

Hormomyces paridiphilus M. Zang & S.L. Wang, Acta bot. Yunn. 19: 324. 1997.

This species was reported as an inhabitant of tubers of *Paris polyphylla* var. *yunnanensis* (*Melanthiaceae*) in China (Zang & Wang 1997). The holotype consists of fragments of uncertain composition overgrown by a dry, greyish oidioid-dendron-like hyphomycete with chains of light brown, subglobose, finely ornamented conidia $(2-2.5-3(3.5) \times (1.5-2-2.5) \mu\text{m}$ (mean $2.78 \pm 0.4 \times 2.35 \pm 0.1$, $Q 1.18$, $n = 50$). Globose chlamydospore-like structures $(7-9-13(15) \times (5-6-10) \mu\text{m}$ (mean $10.1 \pm 0.6 \times 7.89 \pm 0.4$, $Q 1.3$, $n = 15$) composed of oblong, pale brown cells, $2.4(6.5) \times (1-1.5-3.5) \mu\text{m}$ (mean $3.03 \pm 0.1 \times 2.13 \pm 0.1$, $Q 1.45$) were also observed, growing singly and terminally on hyphae. It seems unlikely that the authors would have misinterpreted the *Oidioidendron* as a *Hormomyces*, and dimensions of the observed conidia do not match the protologue; it seems more likely that the original specimen was overgrown by this mould later on. The species will have to be re-collected to re-evaluate its taxonomy. Until then, the taxon should be considered a *nomen dubium*.

Specimen examined: China, Yunnan, on tubers of *Paris polyphylla* var. *yunnanensis*, 10 Sep. 1995, S.L. Wang (holotype HKAS 30237).

Hormomyces peniophorae P. Roberts, Mycotaxon 63: 214, 1997.

The holotype consisted of twigs covered with basidiomes of *Peniophora lycii*, with inconspicuous gelatinous sporodochia ~1 mm diam forming pale spots on the hymenium. Conidia, conidiophores and haustoria consistent with those described by Roberts (1997) were observed. However, because the conidia are not formed in chains, there is no morphological reason to include this fungus in *Hormomyces*, and the presence of "tremellaceous" haustoria described and mycoparasitic ecology are more suggestive of *Tremellales* than *Cantharellales*. Because the haustoria are terminal, Roberts (1997) suggested a possible relationship of his species with the genus *Sirotrema*. However, the species of *Sirotrema* parasitize hosts in *Rhytismataceae*, and its species have clamp connections and yeast-like asexual morphs (Bandoni 1986). *Hormomyces peniophorae* should

be recollected, cultured and sequenced before changes are proposed to its classification.

Specimens examined: UK, England, Devon, Scadson woods, Torquay, as a mycoparasite of *Peniophora lycii* growing on *Rubus idaeus*, 21 Jan. 1996, P. Roberts (holotype K(M):337706); Slapton woods, mycoparasite of *Peniophora lycii* growing on *Ulmus*, 10 Dec. 1994, P. Roberts (K(M):33198).

Hormomyces pezizoideus Speg., Boln Acad. nac. Cienc. Córdoba 2: 467. 1889.

The holotype specimen (LPS 28379) could not be provided on loan, but with the kindness of the curators, we were able to examine Spegazzini's pencil drawing and a macro photograph of the specimen. These suggest a fungus with reddish sporodochia and branched chains of cuboidal or globose conidia ~1.5–2 μm diam, growing on a bamboo. These limited observations do not allow accurate identification of the species, but the small conidial size and bamboo habitat suggest it is not congeneric with *T. aurantaca*.

DISCUSSION

Our taxonomic and phylogenetic revision of the hyphomycete genus *Hormomyces* provides the evidence for the formal synonymy with *Tulasnella* (*Cantharellales*) proposed by Stalpers *et al.* (2021), and the transfer of the type species *H. aurantacus* to the latter genus. We consider two described species to be synonyms of *T. aurantacus* based on type studies, and one species described in *Hormisciopsis* is considered a synonym based on its protologue. *Tulasnella aurantaca* is a relatively common fungus in temperate North America, conspicuous because of its gelatinous orange to reddish sporodochia and abundant globose to ellipsoidal conidia in branched, acropetal chains. While this combination of characters is diagnostic for *T. aurantaca*, other sporodochial fungi with either Basidiomycetous or Ascomycetous affinities could be confused with *T. aurantaca*. A key to similar genera is provided below. The species grows on rotten wood and occasionally overgrows wood-decaying fungi such as *Fomitopsis betulina* and *Crepidotus* sp. We did not observe microscopic structures that might indicate mycoparasitism, such as haustoria, in any specimens, suggesting that *T. aurantaca* sometimes opportunistically overgrows fleshy basidiomes without parasitizing them. *Hormomyces* species are sometimes recorded from unusual substrates. One specimen of a *Hormomyces* sp. was reported from the palm *Rhopalostylis sapida* (McKenzie *et al.* 2004). A single strain identified as *H. aurantacus* was isolated from the surface of a hibernating bat (*Myotis septentrionalis*) collected in a cave in New Brunswick, Canada (Vanderwolf *et al.* 2013). We did not examine either of these collections.

As described above, sporodochial colour is variable in this species. Our interpretation is that differences in colours of sporodochia constitute infraspecific variation related to age or environmental factors, and there are presently no correlations noted with other characters. In several other fungi, pigmentation is influenced by abiotic factors such as light, (e.g. Yu & Fischer 2018). Other sporodochial hyphomycetes, such as *Clonostachys rosea*, are well-known to produce conidiomata with variable colouration, ranging from yellow to orange, pink to red (Schroers

et al. 1999), with a striking bluish green colour defining *C. rosea* f. *catenulata* (Schroers 2001). Variation and overlapping characters in basidiome colour or micromorphology are also problems for the identification of basidiomes of *Tulasnella* species in the absence of DNA sequencing (Cruz et al. 2014).

When this species was known as *H. aurantacus*, it was frequently considered the asexual morph of *Tremella mesenterica*. However, all nuclear DNA analyses confidently place this species in *Tulasnella* among sequences identified as *T. violea*. The latter species is often considered a synonym of *T. lilacina*, the type species of *Tulasnella*, but this synonymy has not yet been evaluated by DNA sequencing. Sequences attributed to *T. violea* appear to be monophyletic in the ITS phylogenetic tree, with the exception of a single strain that may be a misidentified strain of *T. eichleiriana*. In the 28S tree, the two GenBank sequences labelled as *T. violea* are polyphyletic but the strains sampled are different from those in the ITS trees and it is unclear whether one or both are misidentified. Based on our ITS phylogenetic analysis, *Tulasnella* appears to be divided into two clades with *T. aurantaca* in the same clade, with high support, as all the sequences labelled as *T. violea*. If the synonymy of *T. violea* and *T. lilacina* is eventually confirmed, and the concept of the species is clarified, it is probable that *Hormomyces* would remain a synonym of *Tulasnella*.

Until very recently, only nuclear DNA sequences were available for *Tulasnella* species. Sequences for the nuclear glutamate synthase gene (*C4102*), and mitochondrial ATP synthase (*C14436*) were introduced by Arifin et al. (2020) for species from Australia belonging to a clade within *Tulasnella* that includes orchid root endophytes and possibly *T. aurantaca*. Only a few RNA polymerase subunit I (*RPB1*) and RNA polymerase subunit II (*RPB2*) sequences are available (Matheny et al. 2006, Moncalvo et al. 2006). A phylogenetically robust generic concept and any subdivision of *Tulasnella* into segregate genera will require the generation of additional sequences covering the entire family, possibly using *C4102* and *C14436*. Even if such studies justify splitting *Tulasnella* into segregate genera, it is probable that *Hormomyces* would stay nested within any narrower concept

of the genus. Understanding the phylogenetic structure and species concepts in the core clade of *Tulasnella* around the type species clearly requires increased sampling and critical study. The inclusion of strictly asexual species in such analyses should provide additional phylogenetic and morphological characters and perhaps some clarity as future revisions proceed. The eventual taxonomic fate of *Epulorhiza* is less certain. The type species, *E. repens* (= *Tulasnella calospora*), and the other sequenced species of *Epulorhiza* are in the second ITS clade within *Tulasnella*, but this grouping is less evident in the 28S phylogeny. Therefore, it is possible that *Epulorhiza* could be reinstated if *Tulasnella* were divided.

Tulasnella species form mycorrhizae with several types of plants, especially several genera of orchids (Currah et al. 1997, Dearnaley et al. 2012, Almeida et al. 2014, Linde et al. 2017, Oberwinkler et al. 2017, Fujimori et al. 2019, Arifin et al. 2020), and liverworts (Preußing et al. 2009). Using mitochondrial 28S, Almeida et al. (2014) showed that the species formerly included in the asexual genus *Epulorhiza*, namely *T. amonilioides*, *T. epiphytica*, *T. albertaensis* and *T. anatolica*, do not form their own distinct clade, but are distributed across *Tulasnella*. In their phylogenetic analysis, strains identified as the type species of *Epulorhiza*, *E. repens* or its sexual morph *T. calospora*, are polyphyletic, and most of the clades represent unnamed endophytes isolated from orchid roots. We wonder whether *T. aurantacus* may also be able to infect orchid roots because closely related species such as *T. prima*, *T. secunda*, *T. warcupii*, *T. australiensis*, *T. rosea* and several others occur in that niche (Linde et al. 2017, Arifin et al. 2020). *Cantharellales* are known for accelerated evolution of their nuclear DNA loci (Moncalvo et al. 2006) and the mutations in *T. aurantaca* at the binding sites for the universal nuclear DNA primers, ITS4, ITS5 and LROR, may explain why it has not been detected in environmental samples. Different mutations in ITS primer binding sites of *Tulasnella* species were reported by Cruz et al. (2011), who designed specific primers different than those we designed for our study. Our modified primers for *T. aurantacus* (Table 3) may be interesting to try for DNA surveys of orchid mycorrhizae or for the detection of related cryptic species.

Key to genera of hyphomycetes similar to *Tulasnella aurantiaca* (A = Ascomycota, B = Basidiomycota, U = unknown).

- | | |
|--|---------------------------------------|
| 1. Clamp connections present on hyphae | <i>Heteromycophaga</i> (B) |
| 1'. Clamp connections absent on hyphae | 2 |
| 2. No sporodochia produced | 3 |
| 2'. Sporodochia produced | 5 |
| 3. Spore dematiaceous | <i>Phaeomonilia</i> (A) |
| 3. Spore hyaline | 4 |
| 4. Producing large, inflated conidiogenous cells, often on <i>Eucalyptus</i> | <i>Quambalaria</i> (B) |
| 4'. Conidia arising from simple hyphae conidia in long branched chains, only known from cultures derived from ascospores | <i>Chaenothecopsis haematopus</i> (A) |
| 5. Conidia not in chains | 6 |
| 5'. Conidia in chains that may be branched | 8 |
| 6. Sporodochia cup-shaped, conidiogenous cells not in chains | <i>Ditangium</i> (B) |
| 6'. Sporodochia not cup-shaped, conidiogenous cells in chains | 7 |
| 7. Conidiogenous cells clavate | <i>Algonquinia</i> (U) |
| 7'. Conidiogenous cells cuneiform | <i>Catenocuneiphora</i> (U) |

| | |
|--|---|
| 8. Sporodochia gelat nous or slimy | 9 |
| 8'. Sporodochia dry | 14 |
| 9. Conidia 1-septate | <i>Dacrymyces st llatus</i> (B) |
| 9'. Conidia aseptate | 10 |
| 10. On orchid roots, liverworts or slime f uxes on trees | 11 |
| 10'. In beetle galleries, or on rot en wood or mushrooms | 12 |
| 11. Endophyt c on orchid roots, or liverworts | <i>Tulasnella</i> (excluding <i>T. aurant aca</i>)(B) |
| 11'. On slime f uxes on living trees | <i>Oosporidium</i> (A) |
| 12. In beetle galleries | <i>Raf aelea</i> (A) |
| 12'. On rot en wood, lit er or rot en mushrooms | 13 |
| 13. Conidia cylindrical | <i>Calloria</i> (A) |
| 13'. Conidia mostly ellipsoidal | <i>Tulasnella aurant aca</i> (B) |
| 14. Conidia yellow and with a very thick cell wall | <i>Sphaerosporium</i> (A) |
| 14'. Conidia not as above | 15 |
| 15. Parasit c on fresh fruits | <i>Monilia</i> (A) |
| 15'. Not parasit c of fresh fruits | 16 |
| 16. Conidia cuneiform | <i>Hyaloscypha</i> (asexual morphs formerly included in <i>Pseudaegerita</i>)(A) |
| 16'. Conidia ellipsoidal to fusiform, not cuneiform | 17 |
| 17. Conidia hyaline | <i>Cylindrium</i> (A) |
| 17'. Conidia pigmented..... | <i>Hornsmania</i> (U) |

ACKNOWLEDGMENTS

We are grateful to Dr. M. Smith from the Carleton University for his mentorship to JM during this project. We thank Gwenaël Carter, Bibianne Fort n, P. Brandon Matheny and Danny Newman for collect ng fresh specimens, and are indebted to the curators of BPI, DAOM, H, HKAS, and K(M) for lending the specimens used in this study. We are grateful to Dr. S. Redhead for advice on taxonomic and nomenclatural matters, Amy Rossmann and colleagues for helping us resolve the nomenclatural status of *Hormomyces* and *Epulorhiza*, Franck Stefani for assistance with fluorescence microscopy and the Ottawa Research and Development Centre Molecular Technologies Laboratory for DNA sequencing services. We appreciate the efforts of Jorge Chayle, Hugo Madrid and Francisco Kuhar to help us clarify the characters of *H. pezizoideus*. Thanks to Ms. D. Castronovo, Harvard University Faculty of Arts & Sciences, for providing a clean copy of Bonorden's original illustration of *H. aurant acus*.

Conflict of interest: The authors declare that there is no conflict of interest.

REFERENCES

Almeida PRM, van den Berg C, Góes-Neto A (2014). *Epulorhiza amonilloides* sp. nov.: A new species of orchid mycorrhiza from Brazil. *NeoDiversity* 7: 1–10.

Arif n A, May TW, Linde CC (2020). New species of *Tulasnella* associated with Australian terrestrial orchids in the *Cryptostylidinae* and *Drakaeinae*. *Mycologia* DOI: 10.1080/00275514.2020.1813473

Bandoni RJ, Oberwinkler F (1982). *St Ibotulasnella*: A new genus in the

Tulasnellaceae. *Canadian Journal of Botany* 60: 1875–1879.

Bandoni RJ (1986). *Sirotrema*: A new genus in the *Tremellaceae*. *Canadian Journal of Botany* 64: 668–676.

Binder M, Hibbet DS, Larsson KH, et al. (2005). The phylogenetic distribution of resupinate forms across the major clades of mushroom-forming fungi (*Homobasidiomycetes*). *Systematics and Biodiversity* 3: 113–157.

Bonorden HF (1851). *Handbuch der allgemeinen Mykologie*. Schweizerbart'sche Verlagshandlung, Stuttgart, Germany.

Bresadola G (1932). Iconographia mycologica. *Società Botanica Italiana* 23: 1100–1150.

Buyck B, Kauf F, Eyssart er G, et al. (2014). A mult locus phylogeny for worldwide *Cantharellus* (*Cantharellales*, *Agaricomycetes*). *Fungal Diversity* 64: 101–121.

Buyck B, Antonin V, Chakraborty D, et al. (2018). *Cantharellus* sect. *Amethyst ni* in Asia. *Mycological Progress* 17: 917–924.

Celio GJ, Padamsee M, Dent nger BTM, et al. (2006). Assembling the fungal tree of life: constructing the structural and biochemical database. *Mycologia* 98: 850–859.

Cheewangkoon R, Groenewald JZ, Summerell BA, et al. (2009). *Myrtaceae*, a cache of fungal biodiversity. *Persoonia* 23: 55–85.

Crous PW, Wingfeld MJ, Guarro J, et al. (2015). Fungal Planet description sheets: 320–370. *Persoonia* 34: 167–266.

Cruz D, Suárez JP, Kot le I, et al. (2011). Def ning species in *Tulasnella* by correlat ng morphology and nrDNA ITS-5.8S sequence data of basidiomata from a tropical Andean forest. *Mycological Progress* 10: 229–238.

Cruz D, Suárez JP, Kot le I, et al. (2014). Crypt c species revealed by molecular phylogenetic analysis of sequences obtained from basidiomata of *Tulasnella*. *Mycologia* 106: 708–722.

Currah RS, Smreciu EA, Hambleton S (1990). Mycorrhizae and

- mycorrhizal fungi of boreal species of *Platanthera* and *Coeloglossum* (Orchidaceae). *Canadian Journal of Botany* **68**: 1171–1181.
- Currah RS, Zetler LW, McInnis TM (1997). *Epulorhiza inquilina* sp. nov. from *Platanthera* (Orchidaceae) and a key to *Epulorhiza* species. *Mycotaxon* **61**: 335–342.
- Darriba D, Taboada GL, Doallo R, et al. (2012). jModelTest 2: More models, new heuristics and parallel computing. *Nature Methods* **9**: 772.
- Dearnaley, JWD, Martos, F, Selosse, MA (2012). Orchid mycorrhizas: molecular ecology, physiology, evolution and conservation aspects. In: *Fungal Associations* (B. Hock, ed.) Springer-Verlag, Berlin, Germany: 207–230
- Diederich P, Lawrey JD, Capdet M, et al. (2014). New lichen-associated bulbil-forming species of *Cantharellales* (Basidiomycetes). *The Lichenologist* **46**: 333–347.
- Donk MA (1962). The generic names proposed for *Hymenomycetes*. — XII. Deuteromycetes. *Taxon* **11**: 75–104.
- Donk MA (1966). Checklist of European hymenomycetous Heterobasidiales. *Persoonia* **4**: 145–335.
- Edgar RC (2004). MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research* **32**: 1792–1797.
- Feng B, Wang XH, Ratkowsky D, et al. (2016). Multi-locus phylogenetic analyses reveal unexpected abundant diversity and significant disjunct distribution pattern of the Hedgehog Mushrooms (*Hydnum* L.). *Scientific Reports* **6**: 25586.
- Fenwick GA (1995). *Tremella mesenterica* in laboratory culture. *Mycologist* **9**: 124–126.
- Foltz MJ, Perez KE, Volk TJ (2013). Molecular phylogeny and morphology reveal three new species of *Cantharellus* within 20 m of one another in western Wisconsin, USA. *Mycologia* **105**: 447–461.
- Fujimori S, Abe JP, Okane I, et al. (2019). Three new species in the genus *Tulasnella* isolated from orchid mycorrhiza of *Spiranthes sinensis* var. *amoena* (Orchidaceae). *Mycoscience* **60**: 71–81.
- Gams W, van der Aa HA, van der Plaats-Niterink AJ, et al. (1987). *CBS Course of Mycology*, 3rd ed. Baarn and Delft, the Netherlands. Centraalbureau voor Schimmelcultures.
- González V, Salazar O, Julián MC, et al. (2002). *Ceratobasidium albasitensis*: A new *Rhizoctonia*-like fungus isolated in Spain. *Persoonia* **17**: 601–614.
- Guindon S, Gascuel O (2003). A simple, fast and accurate method to estimate large phylogenies by maximum-likelihood. *Systematic Biology* **52**: 696–704.
- Guindon S, Dufayard JF, Lefort V, et al. (2010). New algorithms and methods to estimate maximum-likelihood phylogenies: Assessing the performance of PhyML 3.0. *Systematic Biology* **59**: 307–321.
- Hibbet DS, Luz-Beatrice G, Donoghue MJ (2000). Evolutionary instability of ectomycorrhizal symbioses in basidiomycetes. *Nature* **407**: 506–508.
- Hibbet DS, Binder M (2002). Evolution of complex fruiting-body morphologies in homobasidiomycetes. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **269**: 1963–1969.
- Hoog GS de, Gerrits van den Ende AHG (1998). Molecular diagnostics of clinical strains of filamentous Basidiomycetes. *Mycoses* **41**: 183–189.
- Hopple JS, Vilgalys R (1999). Phylogenetic relationships in the mushroom genus *Coprinus* and dark-spored allies based on sequence data from the nuclear gene coding for the large ribosomal subunit RNA: Divergent domains, outgroups, and monophyly. *Molecular Phylogenetics and Evolution* **13**: 1–19.
- Idris NA (2010). *Identification of a mycorrhiza isolate from Paphiopedilum barbatum* (Lindl.) Pfutzer from ex situ location and determination of its growth-enhancing capabilities. Thesis. Institute of Science, Universiti Putra Malaysia, Malaysia
- Ingold CT (1984). Patterns of ballistospore germination in *Tilletia*, *Auricularia* and *Tulasnella*. *Transactions of the British Mycological Society* **83**: 583–591.
- Jensen-Vargas E, Marizzi C (2018). DNA barcoding for identification of consumer-relevant fungi sold in New York: A powerful tool for citizen scientists? *Food* **7**: 87.
- Kalchbrenner K, Cooke MC (1880). South African fungi. *Grevillea* **9**: 17–34.
- Karsten PA (1890). Fragmenta mycologica XXXI. *Hedwigia* **29**: 270–273.
- Karsten PA (1892). Fragmenta mycologica XXXVIII. *Hedwigia* **31**: 292–296.
- Kennedy LL (1958). The genus *Dacrymyces*. *Mycologia* **50**: 896–915.
- Kornerup A, Wanscher JH (1978). *Methuen Handbook of Colour*. 3rd ed. Eyre Methuen, London, UK.
- Krause C, Garnica S, Bauer R, et al. (2011). *Aneuraceae* (Metzgeriales) and tulasnelloid fungi (Basidiomycota): A model for early steps in fungal symbiosis. *Fungal Biology* **115**: 839–851.
- Kurtzman CP (2011). *Protomyces* Unger (1833). In: *The Yeasts: A Taxonomic Study*, Volume 2 (Kurtzman CP, Fell JW, Boekhout T, eds). Elsevier, Amsterdam, the Netherlands: 725–731.
- Langer EJ (2001). *Phylogeny of non-gilled and gilled basidiomycetes — DNA sequence inference, ultrastructure and comparative morphology*. PhD thesis. University of Tübingen, Germany.
- Larsson KH, Larsson E, Kõljalg U (2004). High phylogenetic diversity among corticioid homobasidiomycetes. *Mycological Research* **108**: 983–1002.
- Lawrey JD, Binder M, Diederich P, et al. (2007). Phylogenetic diversity of lichen-associated homobasidiomycetes. *Molecular Phylogenetics and Evolution* **44**: 778–789.
- Lefort V, Longueville JE, Gascuel O (2017). SMS: Smart Model Selection in PhyML. *Molecular Biology and Evolution* **34**: 2422–2424.
- Linde CC, Phillips RD, Crisp MD, et al. (2013). Congruent species delineation of *Tulasnella* using multiple loci and methods. *New Phytologist* **201**: 6–12.
- Linde CC, May TW, Phillips RD, et al. (2017). New species of *Tulasnella* associated with terrestrial orchids in Australia. *IMA Fungus* **8**: 27–47.
- Lloyd CG (1916). *Hormomyces aurantiacus*. *Mycological Writings* **5**: 170.
- Matheny BP, Gossman JA, Zalar P, et al. (2006). Resolving the phylogenetic position of the *Wallemiomycetes*: An enigmatic major lineage of *Basidiomycota*. *Canadian Journal of Botany* **84**: 1794–1805.
- McCormick MK, Whigham DF, O’Neil J (2004). Mycorrhizal diversity in photosynthetic terrestrial orchids. *New Phytologist* **163**: 425–438.
- McKenzie EHC, Buchanan PK, Johnston PR (2004). Checklist of fungi on Nikau palm (*Rhopalostylis sapida* and *R. baueri* var. *cheesemani*), in New Zealand. *New Zealand Journal of Botany* **42**: 335–355.
- McNabb RFR (1969). The identity of *Hypsilocephala*. *Transactions of the British Mycological Society* **52**: 506–509.
- McNabb RFR (1973). Taxonomic studies in the *Dacrymycetaceae* VIII. *Dacrymyces* Nees ex Fries. *New Zealand Journal of Botany* **11**: 461–524.
- Millanes AM, Diederich P, Ekman S, et al. (2011). Phylogeny and character evolution in the jelly fungi (*Tremellomycetes*, *Basidiomycota*, *Fungi*). *Molecular Phylogenetics and Evolution* **61**: 12–28.
- Moncalvo JM, Lutzoni FM, Rehner SA, et al. (2000). Phylogenetic relationships of agaric fungi based on nuclear large subunit ribosomal DNA sequences. *Systematic Biology* **49**: 278–305.
- Moncalvo JM, Nilsson RH, Koster B, et al. (2006). The cantharelloid

- dade: Dealing with incongruent gene trees and phylogenetic reconstruction methods. *Mycologia* **98**: 937–948.
- MyCoPortal (2019). <http://mycoportal.org/portal/index.php>.
- Nilsson RH, Larsson KH, Larsson E, *et al.* (2006). Fruiting body-guided molecular identification of root-tip mantle mycelia provides strong indications of ectomycorrhizal associations in two species of *Sistotrema* (Basidiomycota). *Mycological Research* **110**: 1426–1432.
- NITE Biological Resource Center (NBRC) (2019). <https://www.nite.go.jp/nbrc/catalogue/NBRCCatalogueDetailServlet?ID=NBRC&CAT=00030400>.
- Oberwinkler F, Cruz D, Suárez JP (2017). Chapter 12. Biogeography and ecology of *Tulasnellaceae*. In: *Biogeography of Mycorrhizal Symbiosis* (Tedersoo L, ed). Springer: Switzerland: 237–271.
- Parthibhan S, Ramasubbu R (2020). Mycorrhizal and endophytic fungal association in *Paphiopedilum druryi* (Bedd.) Stein — A strict endemic and critically endangered orchid of the Western Ghats. *Ecological Genetics and Genomics* **16**: 100059.
- Patouillard N (1900). *Essai taxonomique sur les familles et les genres des Hyménomycètes*. Imprimerie et Lithographie L. Dedume, Lon-le Saunier, France.
- Pereira OL, Rollemberg CL, Borges AC, *et al.* (2003). *Epulorhiza epiphytica* sp. nov. isolated from mycorrhizal roots of epiphytic orchids in Brazil. *Mycoscience* **44**: 153–155.
- Pippola E, Kotiranta H (2008). The genus *Tremella* (Basidiomycota, Tremellales) in Finland. *Annales Botanici Fennici* **45**: 401–434.
- Poli A, Vizzini A, Prigione V, *et al.* (2018). Basidiomycota isolated from the Mediterranean Sea — Phylogeny and putative ecological roles. *Fungal Ecology* **36**: 51–62.
- Preußing M, Nebel M, Oberwinkler F, *et al.* (2010). Diverging diversity patterns in the *Tulasnella* (Basidiomycota, Tulasnellales) mycobionts of *Aneura pinguis* (Marchantophyta, Metzgeriales) from Europe and Ecuador. *Mycorrhiza* **20**: 147–159.
- Rachanarin C, Suwannarach N, Kumla J, *et al.* (2016). A new endophytic fungus, *Tulasnella phuhinrongklaensis* (Cantharellales, Basidiomycota) isolated from roots of the terrestrial orchid, *Phalaenopsis pulcherrima*. *Phytotaxa* **374**: 99–109.
- Raja HA, Miller AN, Pearce CJ, *et al.* (2017). Fungal identification using molecular tools: A primer for the natural products research community. *Journal of Natural Products* **80**: 756–770.
- Rambaut A (2016). FigTree v. 1.4.0. <http://tree.bio.ed.ac.uk/software/figtree/>.
- Raper KB, Thom C (1949). *Manual of the Penicillia*. The Williams & Wilkins Company, Baltimore, United States.
- Roberts P (1994). Globose and ellipsoid-spored *Tulasnella* species from Devon and Surrey, with a key to the genus in Europe. *Mycological Research* **98**: 1431–1452.
- Roberts P (1997). New *Heterobasidiomycetes* from Great Britain. *Mycotaxon* **63**: 195–216.
- Roche SA, Carter RJ, Peakall R, *et al.* (2010). A narrow group of monophyletic *Tulasnella* (Tulasnellaceae) symbiont lineages are associated with multiple species of *Chiloglottis* (Orchidaceae): Implications for orchid diversity. *American Journal of Botany* **97**: 1313–1327.
- Ronquist F, Teslenko M, van der Mark P, *et al.* (2012). MrBayes3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* **61**: 539–542.
- Saccardo PA (1888). *Sylloge Hymenomycetum*, Vol. II. Polyporeae, Hydneae, Thelephoreae, Clavariaeae, Tremellineae. *Sylloge Fungorum* **6**: 1–928.
- Saccardo PA (1916). *Hymeniales (ceterae Agaricaceae, Polyporaceae, Hydnaceae, Thelephoraceae, Tremellaceae)*. *Flora Italica Cryptogama Fungi* **1**: 577–1386.
- Samuels GJ, Ismaiel A, Rosmana A, *et al.* (2012). Vascular Streak Dieback of cacao in Southeast Asia and Melanesia: *In planta* detection of the pathogen and a new taxonomy. *Fungal Biology* **116**: 11–23.
- Schroers H-J, Samuels GJ, Seifert KA, *et al.* (1999). Classification of the mycoparasite *Gliocladium roseum* in *Clonostachys* as *C. rosea*, its relationship to *Bionectria ochroleuca*, and notes on other *Gliocladium*-like fungi. *Mycologia* **91**: 365–385.
- Schroers H-J (2001). A monograph of *Bionectria* (Ascomycota, Hypocreales, Bionectriaceae) and its *Clonostachys* anamorphs. *Studies in Mycology* **46**: 1–211.
- Seifert K, Morgan-Jones G, Gams W, *et al.* (2011). *The Genera of Hyphomycetes*. CBS Biodiversity Series no. 9. 1–997. CBS-KNAW Fungal Biodiversity Centre, Utrecht, Netherlands.
- Shao SC, Buyck B, Hofstetter V, *et al.* (2014). *Cantharellus hygrophorus*, a new species in subgenus *Afrocantharellus* from tropical Southwestern China. *Cryptogamie, Mycologie* **35**: 283–291.
- Shefferson RP, Weiß M, Kull T, *et al.* (2005). High specificity generally characterizes mycorrhizal association in rare lady's slipper orchids, genus *Cypripedium*. *Molecular Ecology* **14**: 613–626.
- Smith MT (1997). *Oosporidium* Stautz. In: *The yeasts — A taxonomic study* (Kurtzman C, Fell JW, eds). Elsevier: Amsterdam, The Netherlands: 598–599.
- Spiegel C (1889). Fungi Puiggarrani. Pugillus 1. *Boletín de la Academia Nacional de Ciencias en Córdoba* **11**: 381–622.
- Stafeu FA, Cowan RS (1976). *Taxonomic literature*. Vol. 1: A–G. 2nd ed. Bohn, Scheltema & Hollemam, Netherlands, Utrecht.
- Stalpers JA, Redhead S, Rossman AY, *et al.* (2021). Competing sexual-aseexual generic names in the *Agaricomycotina* (Basidiomycota), with recommendations for use. *IMA Fungus*: In press.
- Suárez JP, Weiß M, Abele A, *et al.* (2006). Diverse tulasnelloid fungi form mycorrhizas with epiphytic orchids in an Andean cloud forest. *Mycological Research* **110**: 1257–1270.
- Sumst DR (1914). New or interesting fungi. *Mycologia* **6**: 32–36.
- Taylor DL, Bruns TD, Szaro TM, *et al.* (2003). Divergence in mycorrhizal specialization within *Hexaletris spicata* (Orchidaceae), a non-photosynthetic desert orchid. *American Journal of Botany* **90**: 1168–1179.
- Taylor DL, McCormick MK (2008). Internal transcribed spacer primers and sequences for improved characterization of basidiomycetous orchid mycorrhizas. *New Phytologist* **177**: 1020–1033.
- Thiers B (2019). Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/science/ih/>.
- Tubaki K (1976). Cultural and taxonomical studies on *Hormomyces aurantacus*. *Transactions of the Mycological Society of Japan* **17**: 243–247.
- Turland NJ, Wiersema JH, Barrie FR, *et al.* (eds.) (2018). *International Code of Nomenclature for algae, fungi, and plants (Shenzhen Code) adopted by the Nineteenth International Botanical Congress Shenzhen, China, July 2017*. Regnum Vegetabile 159. Koeltz Botanical Books, Glashütten, Germany.
- Van de Put K, Antonissen I (1996). *Tulasnella* uit Vlaanderen. Bijdrage tot de kennis van de Heterobasidiomyceten in Vlaanderen. *Sterbeekia* **17**: 44–69.
- Vandervolf KJ, McAlpine DF, Malloch D, *et al.* (2013). Ectomycota associated with hibernating bats in eastern Canadian caves prior to the emergence of White-nose Syndrome. *Northeastern Naturalist* **20**: 115–130.
- Vasco-Palacios M (2016). *Ectomycorrhizal fungi in Amazonian tropical forests in Colombia*. Thesis. Utrecht University, Netherlands.
- Von Hönel F (1917). Mykologische Fragmente. *Annales Mycologici* **15**: 293–383.

- Vu D, Groenewald M, de Vries M, *et al.* (2019). Large scale generation and analysis of filamentous fungal DNA barcodes boosts coverage for kingdom fungi and reveals thresholds for fungal species and higher taxon delimitation. *Studies in Mycology* **92**: 135–154.
- Weiß M, Bauer R, Sampaio JP, *et al.* (2014). *Tremellomyces* and related groups. In: *The Mycota: Systematics and Evolution* (McLaughlin DJ, Spatafora JW, eds). Springer-Verlag Berlin, Germany: 331–355.
- White TJ, Bruns T, Lee S, *et al.* (1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: *PCR protocols: A guide to the methods and applications* (Innis MA, Gelfand DH, Sninsky JJ, *et al.*, eds). Academic Press, New York, United States: 315–322.
- Wilson N, Hollinger J, *et al.* (2020). Mushroom Observer. <https://mushroomobserver.org/>
- Wu CL, He Y, Yan J, *et al.* (2019). Two new species of *Clavulina* (*Cantharellales*) from southwestern China based on morphological and molecular evidence. *Mycological Progress* **18**: 1071–1078.
- Yu Z, Fischer R (2018). Light sensing and responses in fungi. *Nature Review Biology* **17**: 25–36.
- Zamora JC, Millanes AM, Wedins M, *et al.* (2016). Understanding lichenicolous heterobasidiomycetes: New taxa and reproductive innovations in *Tremella s.l.* *Mycologia* **108**: 381–396.
- Zang M, Wang SL (1997). A new species of the genus *Hormomyces* from China — *Hormomyces paridiphilus*. *Acta Botanica Yunnanica* **19**: 324–324.
- Zelmer CD, Currah RS (1995). *Ceratorhiza pernacetana* and *Epulorhiza calendulina* spp. nov.: Mycorrhizal fungi of terrestrial orchids. *Canadian Journal of Botany* **73**: 1981–1985.

Supplementary Material: <https://fuse-journal.org/>

Figure S1. Mean conidial dimensions (with error bars representing standard error) for all herbarium specimens and cultures examined, with the holotype of *H. fragiforme* represented by orange bars and the holotype of *H. callorioides* represented by green bars.

Figure S2. Lectotype of *H. fragiformis* (A, C, E) and holotype of *H. callorioides* (B, D, F). A, B. Rehydrated sporodochia. C–F. Conidial chains. Scale bars: A, B = 500 µm. C–F = 10 µm.

Figure S3. Conidia and conidial chains. A. *Oosporidium* sp. (DAOM 970823) identified using DNA sequencing. B. Holotype of *Sphaerocolla aurantaca* (H). Both have similar conidial morphology and dimensions, suggesting that *S. aurantaca* may be conspecific with *Oosporidium margaritifera*. Scale bar = 10 µm.

Figure S4. Nuclear staining of hyphae of DAOMC 251988, showing dikaryotic, binucleate hyphae, A, using near-UV light showing the stained nuclei and B with regular light. Scale bar = 20 µm.

Table S1. Species, geographical location, host and herbaria for known type specimens of *Hormomyces* species.

Figure S3

Figure S3. ~~XXXX~~
~~XXXX~~
~~XXXX~~

A. *Oosporidium* ~~XXXX~~

~~XXXX~~

~~XXXX~~

B. ~~XXXX~~

Sphaerocolla
Oosporidium

Figure S4

Figure S4. ~~XXXX~~
~~XXXX~~

Table S1

Hormomyces species.

| Species | XXXX | Host/substrate | Herbarium |
|------------------------|-----------------|---|-----------------|
| XXXX | XXXX | XXXX <i>Pinus</i> and <i>Abies</i> | XXXX |
| XXXX | XXXX | XXXX <i>Quercus</i> | XXXX |
| <i>H. callorioides</i> | XXXX | XXXX | XXXX |
| <i>H. fragiformis</i> | XXXX | XXXX <i>Malus</i> and polypores | XXXX |
| <i>H. paridiphilus</i> | XXXX | <i>Paris polyphylla</i> | XXXX |
| <i>H. peniophorae</i> | XXXX | <i>Peniophora lycii</i> | XXXX |
| <i>H. pezizoideus</i> | XXXX | <i>Bambusaceae</i> | XXXX |